

# Research Report

## Infinitely Renormalizable Diffeomorphisms of the Disk at the Boundary of Chaos

Eleonora Catsigeras

Facultad de Ingenieria IMERL  
C.C. 30, Montevideo, Uruguay

Jean-Marc Gambaudo

IBM Research Division  
T. J. Watson Research Center  
Yorktown Heights, NY 10598

Fernando Moreira

Departamento de Matemáticas  
F.C.U.P., Praça Gomes Teixeira  
4000 Porto, Portugal

### LIMITED DISTRIBUTION NOTICE

This report has been submitted for publication outside of IBM and will probably be copyrighted if accepted for publication. It has been issued as a Research Report for early dissemination of its contents and will be distributed outside of IBM up to one year after the date indicated at the top of this page. In view of the transfer of copyright to the outside publisher, its distribution outside of IBM prior to publication should be limited to peer communications and specific requests. After outside publication, requests should be filled only by reprints or legally obtained copies of the article (e.g., payment of royalties).



Research Division  
Almaden • T.J. Watson • Tokyo • Zurich

# INFINITELY RENORMALIZABLE DIFFEOMORPHISMS OF THE DISK AT THE BOUNDARY OF CHAOS

*Eleonora CATSIGERAS<sup>1</sup>, Jean-Marc GAMBAUDO<sup>2</sup>, and Fernando MOREIRA<sup>3</sup>.*

**Abstract:** We show that, among area contracting embeddings of the 2 disk, infinitely renormalizable maps with a bounded geometry either have positive topological entropy or correspond to a cascade of period doubling.

---

<sup>1</sup> *Facultad de Ingeniería IMERL, C.C. 30, Montevideo, Uruguay.*

<sup>2</sup> *I.N.L.N., 1361, route des lucioles, Sophia-Antipolis, 06560 Valbonne, France.*

<sup>3</sup> *Departamento de Matemáticas, F.C.U.P., Praça Gomes Teireira, 4000 Porto, Portugal.*

## I. Introduction

Let  $f$  be a  $C^1$  map from the unit interval  $I$  into itself and  $(a_n)_{n \geq 0}$  a sequence of integers greater or equal to 2. We say that the map  $f$  is  $(a_n)_{n \geq 0}$ -*infinitely renormalizable* if there exists a sequence of nested intervals  $I \supset I_0(f) \supset I_1(f) \supset \dots \supset I_n(f) \dots$  such that, for each  $n$ :

$$f^i(I_n(f)) \cap I_n(f) = \emptyset, \text{ for } 1 \leq i \leq a_0.a_1 \dots a_n - 1,$$

and

$$f^{a_0.a_1 \dots a_n}(I_n(f)) \subset I_n(f).$$

The intervals  $f^i(I_n)$ , for  $0 \leq i \leq a_0.a_1 \dots a_n - 1$ , are called *the atoms of generation  $n$  of  $f$* .

Maps which satisfy this property but only for a finite sequence  $(a_n)_{m-1 \geq n \geq 0}$  are called  $(a_n)_{m-1 \geq n \geq 0}$ -*renormalizable* or  *$m$ -times renormalizable* when there is no ambiguity.

We say that an infinite renormalisable map is of *bounded combinatorial type* if the sequence  $(a_n)_{n \geq 0}$  is bounded. Notice that, in this case, the sequence  $(a_n)_{n \geq 0}$  has an accumulation point, *i.e.* there is an integer that appears infinitely many times in the sequence.

This type of maps occurs very naturally in one dimensional dynamics: actually for any sequence  $(a_n)_{n \geq 0}$  there exists a value of the parameter  $a$  for which the quadratic map  $x \mapsto 1 - ax^2$  is a  $(a_n)_{n \geq 0}$ -infinitely renormalizable map. Since any continuous map on the interval which possesses a periodic orbit whose period is not a power of 2, has positive topological entropy [BF], it follows easily that the only infinitely renormalizable maps with topological entropy zero are the ones for which each element of the sequence  $(a_n)_{n \geq 0}$  is a power of 2. The aim of this paper is to prove a similar result for area contracting maps of the 2-disk. But, before that, let us emphasize some recent results about infinitely renormalizable maps on the interval.

To a  $(a_n)_{n \geq 0}$ -infinitely renormalizable map  $f$ , we can associate another map, that we call *the renormalized map* of  $f$ , denote by  $\mathcal{R}(f)$ , and define by:

$$\mathcal{R}(f) = \xi^{-1}(f) \circ f^{a_0} \circ \xi(f),$$

where  $\xi(f)$  is an affine scaling which maps  $I$  onto  $I_0(f)$ .

The renormalized map  $\mathcal{R}(f)$  is again a  $(b_n)_{n \geq 0}$ -infinitely renormalizable map with  $b_n = a_{n+1}$  for all  $n$ , and its corresponding sequence of nested intervals is given by:

$$\mathcal{I}_n(\mathcal{R}(f)) = \xi^{-1}(f)(\mathcal{I}_{n+1}(f)) \text{ for all } n \geq 0.$$

From the independent discovery and explanation by Coullet and Tresser ([CT], [TC]) and Feigenbaum [Fe] in 1978, that infinitely renormalizable maps exhibit universal geometrical behaviors, to the culminating work of Sullivan [Su] in 1992, a huge amount of works, both numerical and theoretical, has been done in this field. It is not our intention to give here a panorama of the actual knowledge. For this purpose, we refer the reader to [MS] and the references quoted therein. We just want to focus on a key point of Sullivan's theory, often referred to as "real bounds", that we present here, for sake of simplicity, in a very weakened form:

Consider a smooth map,  $f$ , with a quadratic singularity: more precisely we consider the set  $\mathcal{U}^{1+1}$  of the maps  $f : [0, 1] \rightarrow [0, 1]$  that can be written as

$$f = \phi \circ Q \circ \psi,$$

where  $\psi : [0, 1] \rightarrow [\psi(0), 1]$  is an orientation preserving diffeomorphism such that  $\psi(0)$  is in  $(-1, 0)$ ,  $Q : [\psi(0), 1] \rightarrow [0, 1]$  is the quadratic map  $Q(x) = x^2$ ,  $\phi : [0, 1] \rightarrow [0, 1]$  is an orientation reversing diffeomorphism, and the two maps  $\phi$  and  $\psi$  are  $C^{1+1}$ , that is to say,  $C^1$  and the derivatives satisfy a Lipschitz condition. Sullivan proves the following "beau" theorem:

**Theorem 1.1 [Su]:** *Let  $f \in \mathcal{U}^{1+1}$  be an infinitely renormalizable map with combinatorial type bounded by  $N$ . Then, for all  $n \geq 0$ :*

- 1- *The renormalized maps  $\mathcal{R}^n(f)$  belong to  $\mathcal{U}^{1+1}$ , their  $C^1$ -norm and the Lipschitz constants are bounded by a constant which depends only on  $f$ .*
- 2- *There exist two constants  $a_f$  and  $b_f$  which depend only on  $f$  such that, if  $\mathcal{I}$  is an atom of the generation  $m$  of  $\mathcal{R}^n(f)$  and  $\mathcal{J} \subset \mathcal{I}$  is an atom of the generation  $m + 1$ , then,  $0 < a_f \leq |\mathcal{J}|/|\mathcal{I}| \leq b_f < 1$  (where  $|\cdot|$  stands for the diameter).*

*All these bounds are "beau" (bounded and eventually universally (bounded)), that is to say, that for  $n$  big enough, these bounds can be chosen so that they depend only on  $N$ .*



In the same way as we did in the one-dimensional case, we can define infinitely renormalizable maps in dimension 2. We say that a  $C^1$  map,  $f$ , of the 2-disk  $D^2$  into itself is  $(a_n)_{n \geq 0}$ -infinitely renormalizable if there exists a sequence of nested differentiable disks  $D^2 \supset \mathcal{D}_0 \supset \mathcal{D}_1 \supset \dots \supset \mathcal{D}_n \dots$  such that, for each  $n$ :

$$f^i(\mathcal{D}_n) \cap \mathcal{D}_n = \emptyset, \text{ for } 1 \leq i \leq a_0.a_1 \dots a_n - 1,$$

and

$$f^{a_0.a_1 \dots a_n}(\mathcal{D}_n) \subset \mathcal{D}_n.$$

The sets  $f^i(\mathcal{D}_n)$ , for  $0 \leq i \leq a_0.a_1 \dots a_n - 1$ , are called *the atoms of generation  $n$  of  $f$* . They are not necessarily disks.

In dimension 2, infinitely renormalizable maps are also frequently observed. For instance, they appear naturally in the infinitely dissipative situation for a map  $(x, y) \mapsto (g(x), 0)$ , where  $g$  is an infinitely renormalisable map on the interval, and also in the area preserving case of a map exhibiting resonant islands.

To a  $(a_n)_{n \geq 0}$ -infinitely renormalizable map  $f$  of the 2-disk, we can again associate a renormalized map of  $f$ , defined by:

$$\mathcal{R}(f) = \xi^{-1}(f) \circ f^{a_0} \circ \xi(f),$$

where  $\xi(f)$  is a  $C^1$  scaling which maps  $D^2$  onto  $\mathcal{D}_0(f)$ .

The renormalized map  $\mathcal{R}(f)$  is a  $(b_n)_{n \geq 0}$ -infinitely renormalizable map with  $b_n = a_{n+1}$  for all  $n$ , and its corresponding sequence of nested disks is given by:

$$\mathcal{D}_n(\mathcal{R}(f)) = \xi^{-1}(f)(\mathcal{D}_{n+1}(f)) \text{ for all } n \geq 0.$$

**Definition:** By analogy with the one dimensional case, we say that a  $C^{1+1}$  infinitely renormalizable map of the 2-disk, has a *bounded geometry* if it satisfies the following two conditions:

- 1- For all  $n \geq 0$ , the renormalized maps  $\mathcal{R}^n(f)$ , the scaling maps  $\xi(\mathcal{R}^n(f))$ , and their inverse  $\xi(\mathcal{R}^n(f))^{-1}$ , are  $C^{1+1}$  and their  $C^1$  norm and their Lipschitz constants are bounded by a constant  $K_f$  which only depends on  $f$ .
- 2- There exist constants  $0 < a_f < b_f < 1$  which depends only on  $f$  such that, for all  $n \geq 0$ , if  $\mathcal{I}$  is an atom of the generation  $m$  of  $\mathcal{R}^n(f)$  and  $\mathcal{J} \subset \mathcal{I}$  is an atom of the generation  $m+1$ , then,  $a_f \leq |\mathcal{J}|/|\mathcal{I}| \leq b_f$  ( where  $|\cdot|$  stands for the diameter).

**Remark 1:** These conditions force  $f$  to have a bounded combinatorial type.

**Remark 2:** To have a bounded geometry is a very strong assumption. The above theorem states that an infinitely renormalizable map  $f \in \mathcal{U}^{1+1}$  with bounded combinatorial type, satisfies these assumptions. In fact, this is not known for the other one-dimensional maps (see [Hu]) and obviously not for two-dimensional maps.

Unlike in dimension 1, for any sequence  $(a_n)_{n \geq 0}$ , we can find  $(a_n)_{n \geq 0}$ -infinitely renormalizable  $C^2$  diffeomorphisms of the 2-disk with topological entropy zero; moreover if the sequence  $(a_n)_{n \geq 0}$  is bounded, these maps can be constructed with a bounded geometry [GT2].

However, for an area contracting map of the 2-disk the situation seems to be much more rigid. On one hand, the only known examples of area contracting infinitely renormalizable embeddings with topological entropy zero are such that the sequence  $(a_n)_{n \geq 0}$  is a sequence of powers of 2 [GST]. Actually, these maps are the only known area contracting embeddings of the 2-disk with topological entropy zero, that can be transformed by an arbitrary small  $C^1$ -perturbation into maps with positive topological entropy. On the other hand, there are some numerical evidences that show that for the Hénon model, maps which belong to the  $C^1$ -boundary of positive topological entropy, are geometrically bounded infinitely renormalizable maps of the 2-disk such that the sequence  $(a_n)_{n \geq 0}$  is eventually a sequence of powers of 2 [GT1].

The central result of this paper may be seen as a step towards an explanation of these numerical evidences and can be stated as follows:

**Theorem 1.2:** *Let  $f$  be a  $(a_n)_{n \geq 0}$ -infinitely renormalizable map of the 2-disk with a bounded geometry, which contracts uniformly the area. Then:*

- *either, the topological entropy of  $f$  is positive,*
- *or, eventually the sequence  $(a_n)_{n \geq 0}$  is a sequence of powers of 2.*

### III. Proof of the Theorem

Let us start with some notations. For any positive  $K$ , we denote by  $\mathcal{U}(K)$  the set of  $C^{1+1}$  maps from the disk  $\mathbf{D}^2$  into itself, whose derivatives are Lipschitz, with Lipschitz constant smaller than  $K$ . Thanks to the Arzela-Ascoli theorem, any sequence of maps in  $\mathcal{U}(K)$  has a converging subsequence in the  $C^1$  topology. All along the proof, we shall frequently make use of this property.

Consider now an  $(a_n)_{n \geq 0}$  infinitely renormalizable map  $f$  of the 2-disk with a bounded geometry. Since the sequence  $(\mathcal{R}^n(f))_{n \geq 0}$  remains in  $\mathcal{U}(K_f)$ , it possesses an accumulation point in  $\mathcal{U}(K_f)$ .

**Lemma 3.1:** *Let  $p_0$  be an accumulation point of the sequence  $(a_n)_{n \geq 0}$ . Then, there is a map  $g_0$  in  $\mathcal{U}(K_f)$  which is an accumulation point of the sequence  $(\mathcal{R}^n(f))_{n \geq 0}$  and which satisfies:*

- (i)  $g_0$  is 1-time renormalizable, more precisely there exists a differentiable disk  $\mathcal{D}_0(g_0) \subset \mathbf{D}^2$  such that  $\mathcal{D}_0(g_0), g_0(\mathcal{D}_0(g_0)), \dots, g_0^{p_0-1}(\mathcal{D}_0(g_0))$  are disjoint and  $g_0^{p_0}(\mathcal{D}_0(g_0)) \subset \mathcal{D}_0(g_0)$ .
- (ii) Every atom  $\mathcal{J}$  of the first generation of  $g_0$  satisfies  $|\mathcal{J}| \leq 2b_f$  (where  $b_f$  is the bound given in the above definition).

**Proof:** Let  $p_0$  be an accumulation point of the sequence  $(a_n)_{n \geq 0}$ . Then, there exists a subsequence  $(a_{\phi(n)})_{n \geq 0}$  which is constant and equal to  $p_0$ . From the sequences  $\mathcal{R}^{\phi(n)}(f)$  and  $\xi(\mathcal{R}^{\phi(n)}(f))$ , we can extract subsequences  $\mathcal{R}^{\psi(n)}(f)$  and  $\xi(\mathcal{R}^{\psi(n)}(f))$  which respectively converge to maps  $g_0$  and  $\xi_0$  in  $\mathcal{U}(K_f)$ .

For each  $n \geq 0$ , we have:

$$\mathcal{D}_0(\mathcal{R}^{\psi(n)}(f)) = \xi(\mathcal{R}^{\psi(n)}(f))(\mathbf{D}^2).$$

Since the maps  $\mathcal{R}^{\psi(n)}(f)$  are 1-time renormalizable, we get:

$$(\mathcal{R}^{\psi(n)}(f))^i(\mathcal{D}_0(\mathcal{R}^{\psi(n)}(f))) \cap \mathcal{D}_0(\mathcal{R}^{\psi(n)}(f)) = \emptyset \text{ for } 1 \leq i \leq p_0 - 1,$$

and

$$(\mathcal{R}^{\psi(n)}(f))^{p_0}(\mathcal{D}_0(\mathcal{R}^{\psi(n)}(f))) \subset \mathcal{D}_0(\mathcal{R}^{\psi(n)}(f)).$$



By setting

$$\mathcal{D}_0(g_0) = \xi_0(\mathbf{D}^2),$$

we get by continuity:

$$(g_0^i(\mathcal{D}_0(g_0)) \cap \mathcal{D}_0(g_0) = \emptyset \text{ for } 1 \leq i \leq p_0 - 1,$$

and

$$g_0^{p_0}(\mathcal{D}_0(\mathcal{D}_0(g_0))) \subset \mathcal{D}_0(g_0).$$

The fact that  $\mathcal{D}_0(g_0)$  is a differentiable disk comes from the uniform estimates on the norm of the derivative of the inverse scaling functions.

Since  $f$  has bounded geometry, we know that, for each  $n \geq 0$  and for each atom  $J$  of the first generation of  $\mathcal{R}^n(f)$ , we have:

$$|J| \leq 2b_f.$$

By continuity we get the same estimate for  $g_0$ .

(q.e.d.)

Lemma 3.1 is actually the first step of an inductive process:

**Lemma 3.2:** *Let  $p_0$  be an accumulation point of the sequence  $(a_n)_{n \geq 0}$ . Then, there exists a sequence of maps  $(g_l)_{l \geq 0}$  in  $\mathcal{U}(K_f)$  which are accumulation points of the sequence  $(\mathcal{R}^n(f))_{n \geq 0}$  and which satisfy, for each  $l \geq 0$ :*

(i)  $g_l$  is  $l+1$  times renormalizable. More precisely, there exists a sequence  $(a_{l,n})_{l \geq n \geq 0}$  with  $a_{l,l} = p_0$ , such that  $g_l$  is  $(a_{l,n})_{l \geq n \geq 0}$ -renormalizable.

(ii)  $\mathcal{R}(g_l) = g_{l+1}$ .

(iii) Every atom  $\mathcal{J}$  of the  $n^{\text{th}}$  generation of  $g_l$ ,  $0 \leq n \leq l+1$ , satisfies  $|\mathcal{J}| \leq 2b_f^n$ .

**Proof:** Let  $p_0$  be an accumulation point of the sequence  $(a_n)_{n \geq 0}$ . There exists a subsequence  $(a_{\phi(n)})_{n \geq 0}$  which is constant and equal to  $p_0$ . The subsequence  $(a_{\phi(n)-1})_{n \geq 0}$  has also an accumulation point, say  $p_1$ . By iterating this process  $l$  times, we can find a subsequence  $(a_{\phi_l(n)})_{n \geq 0}$  which is such that:

- $a_{\phi_l(n)}$  is constant and equal to  $p_l$ ,
- $a_{\phi_l(n)+1}$  is constant and equal to  $p_{l-1}$ ,
- ...



$a_{\phi_l(n)+l}$  is constant and equal to  $p_0$ .

By a diagonal process, we can extract from the sequences  $\mathcal{R}^{\phi_l(n)}(f)$  and  $\xi(\mathcal{R}^{\phi_l(n)}(f))$ , subsequences  $\mathcal{R}^{\psi_l(n)}(f)$  and  $\xi(\mathcal{R}^{\psi_l(n)}(f))$  which respectively converge to maps  $g_l$  and  $\xi_l$  in  $\mathcal{U}(K_f)$  and  $g_l$  is such that  $\mathcal{R}(g_l) = g_{l-1}$ . From this point, we can use the same techniques as in the proof of Lemma 3.1, to terminate the proof of Lemma 3.2.

(q.e.d.)

**Lemma 3.3:** *For each  $l \geq 0$ , there exist an atom  $J_l$ , of the  $l^{th}$  generation of  $g_l$  and a point  $x_l$  in  $J_l$  such that  $\|dg_l(x_l)\| \geq 1$ .*

**Proof 3.3:** We know that

$$g_l^{p_1 \dots p_l}(\mathcal{D}_l(g_l)) \subset \mathcal{D}_l(g_l),$$

where

$$\mathcal{D}_l(g_l) = \xi_0 \dots \xi_{l-1}(\mathbf{D}^2),$$

and that  $g_l^{p_1 \dots p_l}$  possesses in  $\mathcal{D}_l(g_l)$  a periodic orbit with period  $p_0$ .

It follows that there exists a point  $y_l$  in  $\mathcal{D}_l(g_l)$  such that

$$\|dg_l^{p_1 \dots p_l}(y_l)\| \geq 1.$$

Consequently, in one of the  $p_1 \dots p_l - 1$  first images of  $\mathcal{D}_l(g_l)$ , that is to say, in an atom  $J_l$ , of the  $l^{th}$  generation of  $g_l$  there is a point  $x_l$ , image of  $y_l$  by some iterate of  $f$ , such that  $\|dg_l(x_l)\| \geq 1$ .

(q.e.d.)

Let us now assume that the map  $f$  contracts uniformly the area, i.e. there exists  $b$  such that  $|\det(df(x))| \leq b < 1$  for all  $x$  in  $\mathbf{D}^2$ . Then we have the following result:

**Lemma 3.4:** *Any accumulation point  $g$  of the sequence  $(\mathcal{R}^n(f))_{n \geq 0}$  is a singular map, i.e.  $\det(dg_0(x)) = 0$  for all  $x \in \mathbf{D}^2$*

**Proof:** We have

$$\mathcal{R}^n(f) = \xi^{-1}(\mathcal{R}^{n-1}(f)) \circ \dots \circ \xi^{-1}(f) \circ f^{a_0 \dots a_{n-1}} \circ \xi(f) \circ \dots \circ \xi(\mathcal{R}^{n-1}(f)).$$

Since for any linear map  $A$  in finite dimension, we have  $|\det A| \leq \|A\|^2$ , it follows that:

$$|\det d\mathcal{R}^n(f)(x)| \leq K_f^{4n} b^{a_0 \dots a_{n-1}}.$$

and this quantity goes to 0 when  $n$  goes to  $\infty$ . Thus, by continuity, any accumulation point of the sequence  $(\mathcal{R}^n(f))_{n \geq 0}$  is a singular map.

(q.e.d.)

Consider now the sequence  $(g_l, x_l)$  defined in Lemmas 3.1 and 3.2. We can extract from it, a subsequence  $(g_{\theta(l)}, x_{\theta(l)})$  which converges to some  $(g, x)$  where  $g$  is a map in  $\mathcal{U}(K_f)$  which is an accumulation point of the sequence  $(\mathcal{R}^n(f))_{n \geq 0}$ , and  $x$  is an accumulation of the sequence  $(x_l)_{l \geq 0}$ . By continuity, we get  $\|dg(x)\| \geq 1$ . It follows that there exist a connected neighbourhood  $\mathcal{V}$  of  $x$  and an integer  $l_0$  such that, for all  $y \in \mathcal{V}$ , and for all  $l \geq l_0$ :

$$(i) \|dg_{\theta(l)}(y)\| \geq 1/2,$$

and

$$(ii) \mathcal{V} \text{ contains } J_{\theta(l)} \text{ (the atom of the } \theta(l)^{\text{th}} \text{ generation of } g_{\theta(l)} \text{ containing } y_{\theta(l)}).$$

Thanks to Lemma 3.4, we know that the maps  $g_l$  and  $g$  are singular:

$$\det(dg_l(y)) = \det(dg(y)) = 0, \forall y \in \mathbf{D}^2.$$

Therefore, for all  $l \geq l_0$ , and for all  $y \in \mathcal{V}$ , the dimension of the kernel of  $Dg_{\theta(l)}$  is 1. Thus,  $\ker Dg_{\theta(l)}$  defines a Lipschitz field of directions on  $\mathcal{V}$ , and consequently a Lipschitz foliation. The image of each leaf is a point because the derivative of  $g_{\theta(l)}$  along the leaves is zero. Consider now a trivialisation of this foliation in  $\mathcal{V}$ . That is to say a map  $\tau : \mathcal{V} \rightarrow \mathbf{R}^2$  which is a Lipschitz homeomorphism onto its image and that maps each leaf of the foliation in  $\mathcal{V}$  into a horizontal line. Recall now that  $g_{\theta(l)}$  maps its atom of the  $\theta(l)^{\text{th}}$  generation,  $J_{\theta(l)}$ , into itself after  $q_{\theta(l)} = p_1 \dots p_{\theta(l)}$  iterations:

$$g_{\theta(l)}^{q_{\theta(l)}}(J_{\theta(l)}) \subset J_{\theta(l)},$$

and that  $g_{\theta(l)}^{q_{\theta(l)}}$  possesses in  $J_{\theta(l)}$ , and thus in  $\mathcal{V}$ , a periodic orbit with period  $p_0$ .

Using the conjugacy by the trivialisation  $\tau$ , the map  $f_{\theta(l)} = \tau \circ g_{\theta(l)}^{q_{\theta(l)}} \circ \tau^{-1} : \tau(\mathcal{V}) \rightarrow \mathbf{R}^2$  reads:

$$f_{\theta(l)}(x_1, x_2) = (f_{1, \theta(l)}(x_2), f_{2, \theta(l)}(x_2)),$$

where  $(x_1, x_2)$  are the canonical coordinates in  $\mathbf{R}^2$  and  $f_1$  and  $f_2$  are two continuous maps from some interval to the reals.

Since the map  $g_{\theta(I)}^{q_{\theta(I)}}$  has a periodic orbit with period  $p_0$  in  $\mathcal{V}$ , we get that the map  $f_{2,\theta(I)}(x_2)$  possesses also a periodic orbit with period  $p_0$ .

We are now in a good position to prove our theorem. Assume that we started with a  $(a_n)_{n \geq 0}$ -infinitely renormalizable map  $f$  of the 2-disk with a bounded geometry. Assume also that  $f$  contracts uniformly the area. If the sequence  $(a_n)_{n \geq 0}$  is not eventually a sequence of powers of 2, it has an accumulation point  $p_0$  which is not a power of 2. The construction we made above yields an interval map  $f_{2,\theta(I)}$  which has a periodic orbit whose period  $p_0$  is not a power of 2, and thus has positive topological entropy. This means [Mi] that there exist an interval  $I$  where  $f_{2,\theta(I)}$  is defined, two disjoint subintervals  $I_0 \subset I$  and  $I_1 \subset I$ , and an integer  $n$  such that:

$$f_{2,\theta(I)}^n(I_0) \supset I,$$

and

$$f_{2,\theta(I)}^n(I_1) \supset I.$$

It follows that the map  $f_{\theta(I)}^n$  maps the two horizontal strips  $D_0 = (\mathbf{R} \times I_0) \cap \tau(\mathcal{V})$  and  $D_1 = (\mathbf{R} \times I_1) \cap \tau(\mathcal{V})$  on two lines whose projections on the vertical axis (parallelly to the horizontal one) cover the interval  $I$ .

Consider now a continuous map  $g : \tau(\mathcal{V}) \rightarrow \tau(\mathcal{V})$ . If  $g$  is  $C^0$  close enough to the map  $f_{\theta(I)}$ , it will map the two strips  $D_0$  and  $D_1$  on two strips whose projections on the vertical axis (parallelly to the horizontal one) cover the interval  $I$  and such that  $g^n(\mathbf{R} \times \partial I_0) \cap \tau(\mathcal{V})$  and  $g^n(\mathbf{R} \times \partial I_1) \cap \tau(\mathcal{V})$  do not intersect  $\mathbf{R} \times I$ , where  $\partial I$  stands for the boundary of  $I$ .

It follows that  $g^n$  has an invariant set in  $\tau(\mathcal{V})$  such that  $g^n$ , restricted to this invariant set, is semi-conjugate to the shift on two symbols. Thus  $g^n$ , and consequently  $g$ , have positive topological entropy. Since the map  $g_{\theta(I)}$  is an accumulation point of the sequence  $(\mathcal{R}^n(f))_{n \geq 0}$ , we get that some iterates of maps in this sequence accumulate (in the  $C^1$ -topology) on the map  $g_{\theta(I)}^{q_{\theta(I)}}$ . Thus, their images by the conjugacy by  $\tau$  accumulate, in the  $C^0$ -topology, on  $f_{\theta(I)}$ . Consequently the renormalized maps  $(\mathcal{R}^n(f))_{n \geq 0}$  and then  $f$  have positive topological entropy. This achieves the proof of our theorem.

**Acknowledgements:** The three authors thank the IMPA from Rio de Janeiro, where this work started, and particularly J. Palis for stimulating discussions. J.M.G. thanks IBM T.J. Watson center for its hospitality during the final stage of this work. F.M. has been



partially supported by JNICT and project JNICT/PBIC// C/CEN/1020/92.

### References.

- [BF] R. Bowen and J. Franks, "The periodic points of maps of the disk and the interval," *Topology* **15** (1976) 337-342.
- [CT] P. Coullet and C. Tresser, "Itérations d'endomorphismes et groupe de renormalisation," *J. Phys. C5* (1978) 25-28.
- [GST] J.M. Gambaudo, S. van Strien and C. Tresser, "There exists a  $C^\infty$  Kupka-Smale diffeomorphism on  $\mathbb{R}^2$  with neither sinks nor sources," *Nonlinearity* **2** (1989) 287-304.
- [GT1] J.M. Gambaudo and C. Tresser, "How horseshoes are created," in " **Instabilities and Nonequilibrium Structures III**" E. Tirapegui and W. Zeller Eds. (Reidel; Dordrecht/Boston/Lancaster/Tokyo) (1991).
- [GT2] J.M. Gambaudo and C. Tresser, "Self-similar constructions in smooth dynamics. Smoothness and dimension," *Commun. Math. Phys.* **150** (1992) 45-58.
- [Fe] M. Feigenbaum, "Quantitative universality for a class of non-linear transformations," *J. Stat. Phys.* **19** (1978) 25-52.
- [Hu] J. Hu, "Renormalization, rigidity and universality in bifurcation theory" *Ph. D. Dissertation, CUNY* (1979).
- [MS] W. de Melo and S. van Strien, **One Dimensional Dynamics** (Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, Vol. 25) (Springer-Verlag, Berlin, 1993).
- [Mi] M. Misiurewicz, "Horseshoes for mappings of the interval," *Bull. Acad. Pol. Ser. Sci. Math.* **27** (1979) 167-169.
- [Su] D. Sullivan, "Bounds, Quadratic differentials, and Renormalization Conjectures," in *A.M.S. Centennial Publication, Vol. (Providence, RI)* (1992).
- [TC] C. Tresser and P. Coullet, "Itérations d'endomorphismes et groupe de renormalisation," *C. R. Acad. Sc. Paris* **287A** (1978) 577-580.