

CHAOS AND STABILITY IN A MODEL OF INHIBITORY NEURONAL NETWORK.

Eleonora Catsigeras *

November 28, 2008

Abstract

Abstract. We analyze the dynamics of a deterministic model of inhibitory neuronal networks proving that the discontinuities of the Poincaré map produce a never empty chaotic set, while its continuity pieces produce stable orbits. We classify the systems in three types: the almost everywhere (a.e.) chaotic, the a.e. stable, and the combined systems. The a.e. stable are periodic and chaos appears as bifurcations. We prove that a.e. stable systems exhibit limit cycles, attracting a.e. the orbits.

Keywords: Chaotic sets, limit cycles, piecewise continuous dynamics, neuronal networks.

1. INTRODUCTION

We obtain rigorous mathematical new results on the dynamics of a deterministic abstract discontinuous dynamical system, in a finite but large dimensional phase space. It comes from a non linear model of inhibitory neuronal networks, without delays, composed by equally or different $2 \leq n < +\infty$ pacemaker neurons, evolving according to an autonomous differential equation in the interspike interval times, and interacting among them by synaptical instantaneous currents in the spiking instants.

The phase space is a bounded cube $Q = [-\theta, \theta]^n$ of the vectorial space \mathbb{R}^n , whose points are the n -uples $V = (V_1, \dots, V_n)$ with $-\theta \leq V_i \leq \theta, \forall 1 \leq i \leq n$. We denote

$$I = \{1, 2, \dots, n\}$$

to the set of neurons of the network. The variable V_i is the potential states of the i -th. neuron of the network, and it evolves on time t .

In this model the synapses and the spikes of the neurons, can be produced at any instant t_i : when the potential V_i of at least one of the neurons (say the i -th neuron), reaches θ (the fixed threshold level). t_i is not previously specified: it is the solution of the implicit function $\Phi_i^t = \theta$. We denote as Φ_i^t to the i -th. component of the vectorial flux solution Φ^t of an autonomous differential equation

$$dV/dt = F(V), \forall V \in Q \subset \mathbb{R}^n, F : Q \mapsto \mathbb{R}^n, F \in C^1 \quad (1)$$

This vectorial differential equation determines the potential V_i of each neuron i as a function on time t , only during the interspike regime. The solution $\Phi^t(V^0)$ depends on the initial state

*Instituto de Matemática y Estadística Prof. Rafael Laguardia. Facultad de Ingeniería, Universidad de la República. Uruguay.

$V^0 = (V_1^0, \dots, V_n^0)$. We restrict to the case in which the system is composed by n independent differential equations (i.e. F_i depends only of V_i). Nevertheless the open conclusions we obtain (for instance the result in Theorem 3) holds even if the neurons slightly interact during the interspike intervals (i.e. $\partial F_i / \partial V_j \approx 0$ if $i \neq j$). We assume that

$$F_i > 0, \quad \partial F_i / \partial V_i < 0 \quad \forall i \in I, \quad \partial F_i / \partial V_j = 0 \quad \forall j \neq i \quad (2)$$

and F_i and $\partial F_i / \partial V_i$ are bounded away from zero. The dynamics given by (1) and (2) is that of general pacemaker neurons, with strictly increasing potentials in dissipative regime, providing negative concavity of the free evolution on time, of the potential of each neuron in the interspike time intervals (ISI). This property leads to a contractive Poincaré map (Theorem 3 and Budelli *et al.*[1996]), and produce, when synaptic inhibitory coupling is added, asymptotic cyclic attractors in the phase space (see Theorem 8), but also chaotic orbits (see Theorem 6). The limit cycles exhibit mostly asynchronous spikes, but in periodic patterns, although they could seem irregular to an experimental observer, if the period is very large.

The model verifying (1) and (2) includes the networks of coupled leaky integrators and the relaxation oscillators, for instance if $F_i = -\gamma_i(V_i - \beta_i)$, with constant $0 < \gamma_i, \beta_i$. The mathematical approach, under particular hypothesis, on the integrate and fire and the relaxation oscillator models of n neurons was early analyzed by Mirollo & Strogatz [1990] for homogeneous networks of excitatory neurons. Later Abbott & van Vreeswijk [1993] generalized the results for non homogeneous systems, also with delay times and excitatory interactions. For strongly coupled networks of inhibitory or excitatory neurons Bessloff & Coombes [2000] find synchronous and asynchronous behavior.

From an initial state V^0 , the next spiking instant is

$$\bar{t}(V^0) = \min_{1 \leq i \leq n} t_i(V^0) \quad (3)$$

where $t_i(V^0)$ is the solution of the implicit function

$$\Phi_i^t(V^0) = \theta \Leftrightarrow t = t_i(V^0) \quad (4)$$

At instant \bar{t} maybe more than one neuron reaches the threshold level θ simultaneously (although this occurs with zero Lebesgue probability in the initial state, if the system is inhibitory). We call $J(V^0)$ to that set of neurons that spike first, from the initial state V^0 , i.e.

$$J(V^0) = \{i \in I : t_i(v^0) = \bar{t}(V^0)\} \quad (5)$$

Almost all initial states are such that $J(V^0) = \{i\}$ for a single neuron $i \in I$. The set J of neurons gives spikes at time \bar{t} , i.e. their potentials reset to zero, and from the new initial state W^0 it restart the evolution according to the differential equation (1). Precisely

$$V_i(\bar{t}^-) = \theta \Rightarrow V_i(\bar{t}^+) = W_i(0) = 0 \quad \forall i \in J \quad (6)$$

When spiking, each of the neurons $i \in J$, which are supposed to be inhibitory, produces a sudden synaptic current through its connections to the other neurons $j \notin J$. This synapsis produces a sudden change, of amplitude $-H_{ij} < 0$, in the potential V_j . Precisely:

$$V_j(\bar{t}^-) < \theta, \quad V_j(\bar{t}^+) = W_j(0) = V_j(\bar{t}^-) - \sum_{i \in J} H_{ij} \quad \forall j \notin J \quad (7)$$

In the abstract dynamical system we define the complementary sets C and S , of chaotic and stable future orbits respectively (Section 3). The dynamics of the first return map $\rho : B \mapsto B$

is adequately defined in a Poincaré section B . It will be diffeomorphic to the union of a finite number of $n - 1$ dimensional balls transverse to the flux.

The qualitative rigorous analysis, discretizing the dynamics by means of a Poincaré map, was applied to homogeneous network networks by Mirolo & Strogatz [1990]. Also Budelli *et al.* [1991] and Catsigeras & Budelli [1992] studied the asymptotic future behavior of inhomogeneous two neurons networks, with a Poincaré map in a codimension one section.

On the other hand discretizations on time, are different mathematical approaches and can fit better with computer experiments, in which both time and space are discretized. Cessac [2008] and Césac & Viéville [2008] obtain similar results to those in this paper, using different mathematical models. They also study discrete discontinuous neural networks. Their model, in spite of being similar in the evolution given by equation (1), is intrinsically different to consider the spike instants as predetermined by the observer.

In this paper we prove that the Poincaré map ρ is discontinuous due to synapsis. Although ρ is piecewise contractive, we prove that its discontinuities play the role of chaos generators: due to them *the chaotic set C is never empty*.

The abstract dynamical systems of this model are classified in three types according to their attractors a.e.. Calling m to the $n - 1$ - dimensional Lebesgue probability measure in the Poincaré section B , we define the *a.e. chaotic systems*, for which C has full measure: $m(C) = 1$, $m(S) = 0$; the *a.e. stable systems* for which $m(C) = 0$, $m(S) = 1$, and *the combined systems*, for which $m(C) > 0$, $m(S) > 0$.

Generically, under some additional hypothesis, *the systems are a.e. stable, and the chaotic set appears as a bifurcation* among different stable systems. This kind of bifurcations were studied by Catsigeras, Rovella & Budelli [2008], proving that they exhibit a chaotic Cantor set attractor.

In Theorem 8 of this paper we prove that *the a.e. stable systems exhibit limit cycles attracting Lebesgue almost all the orbits*, and due to them they are a.e. periodic. Nevertheless the period of the attracting limit cycles, and the number of them, are mostly determined by the relation among the inhibitory synaptic interactions. They can a priori be very large, and have few relation with the intrinsic individual periods of the integrate and fire neurons of the network. This last result fits with the numerical experiments in networks of n different integrate and fire neurons. Postnova *et al.*[2007] analyzed the dynamics of computer simulated large networks, and obtained that the system can be driven through different synchronization states, but they are significantly different from the original periodic behavior of the individual cells.

2. THE POINCARÉ SECTION AND THE FIRST RETURN MAP.

In the compact phase space $Q = \{V \in \mathbb{R}^n : -\theta \leq V_i \leq \theta \forall i \in I\}$ we take the following $n - 1$ -dimensional section:

$$B = \bigcup_{k=1}^n \widehat{B}_k \quad \text{where} \quad \widehat{B}_k = \{V \in Q : V_k = 0\}$$

Its topology is that induced by $\widehat{B}_k \subset \mathbb{R}^{n-1}$. B is transversal to the flux Φ^t . In fact, $d\Phi_i^t/dt = F_i(\Phi^t) > 0$ because (2). Due to the definition of the mathematical spike in equation (6), from any initial state $V^0 \in Q$ the system arrives to the Poincaré section B in a finite time $\bar{t}(V^0)$ given by equations (3) and (4). So in particular for $V^0 \in B$, we have defined the first return Poincaré map ρ :

$$\rho : B \mapsto B, \quad \rho(V^0) = W^0 = V^{\bar{t}^+(V^0)} \quad (8)$$

The sequence of interspike intervals (ISI) is given by the evolution of the system through the iterates ρ^p , $p = 1, 2, \dots$ of the return map ρ . The ISI sequence is $\bar{t}(V^0), \bar{t}(\rho(V^0)), \bar{t}(\rho^2(V^0)), \dots, \bar{t}(\rho^p(V^0)), \dots$. If the orbits of the discrete dynamical system by iteration of ρ are attracted to limit cycles or not,

the ISI sequence will be finally periodic or not. If the dynamics of ρ exhibits chaotic attractors, then the ISI sequence also. Even in the periodic case, its period depends of the map ρ , (and also of the initial state if there were many limit cycles). But ρ depends strongly of the matrix of synaptic interactions H_{ij} . Therefore, the network composed by the inhibitory coupling of n oscillators, may have a dynamics which widely differs from the behavior of each isolated neuron.

We iterate the Poincaré map $\rho(V^0) = W^0$, and after each iterate we reset the time to consider the solution $\Phi^t(W^0)$ of the equation (1), from the new initial state $W^0 \in B$. To simplify the notation we will omit the supraindex 0 : from now U, V, W denote points in the Poincaré section.

From equalities (6), (7) and (8) we have the following formula for the Poincaré map:

$$\begin{aligned} \rho_j(V) &= \Phi^{\bar{t}(V)}(V) - \sum_{i \in J(V)} H_{ij} \quad \forall j \notin J(V) \\ \rho_i(V) &= 0 \quad \forall i \in J(V) \end{aligned} \quad (9)$$

Denote $\wp(I)$ to the family of all non empty parts of the set I of neurons. Roughly speaking $\wp(I)$ the collection of all possible “words” of *different* neurons, of any length $\geq 1, \leq n$, without considering the order. For each $J \in \wp(I)$ (say for instance, $J = \{1, 2\}$), define:

$$B_J = \{V \in B : J(V) = J\}$$

In the example $J = \{1, 2\}$, the set $B_{\{1,2\}} \subset B$ is composed by all the initial states in the Poincaré section B for which neurons 1 and 2, and only them, will arrive first and simultaneously to the threshold level.

Remark 1. Due to formulas (9) the Poincaré map ρ is *continuous in each piece B_J , in particular in $B_{\{i\}}$ for all $i \in I$* . The set B_J , for each $J \in \wp(I)$, is called a *continuity piece* of ρ .

From (9), due to the transversality of the flux Φ^t to \widehat{B}_i , ρ transforms homeomorphically $B_{\{i\}} \cap \widehat{B}_k$ onto its image in \widehat{B}_i . So it is an *open* map. Nevertheless it is not globally injective because each point may have preimages in B_k and B_h for $h \neq k$.

The partition \mathcal{P} and the set $\partial\mathcal{P}$ of discontinuities.

By construction, for different subsets $J^1 \neq J^2 \in \wp(I)$: $B_{J^1} \cap B_{J^2} = \emptyset$. Any initial state V in B belongs to one and only one set B_J (exactly to that B_J such that $J(V) = J$). Therefore:

The collection $\{B_J\}_{J \in \wp(I)}$ form a partition \mathcal{P} of the Poincaré section B into the continuity pieces of the return map ρ .

(3) and (4) implies that the set of $V \in B$ such that $\bar{t}(V) = t_i(V) < t_j(V) \quad \forall j \neq i$ is open. In other words it is open the set $B_{\{i\}}$ (i.e. the set of initial states from where the neuron i , and only the neuron i , arrives first to the threshold level). On the other hand B_J has empty interior if $\#J \geq 2$.

We consider only those *not dead* neurons (i.e. they do not remain under the threshold level forever) and rename as I to that set of neurons. If there is only one, then there is not interactive network to study. So let us suppose that $\#I \geq 2$ and then $\{B_i \neq \emptyset\}_{i \in I}$ is a collection of at least two open and non empty sets. As the space B is connected, we conclude that *the topological frontiers of the pieces of the partition \mathcal{P} is not empty*. Define:

$$\partial\mathcal{P} = B \setminus \bigcup_{i=1}^n B_{\{i\}} = \bigcup_{\#J \geq 2} B_J = \bigcup_{i=1}^n \partial B_{\{i\}}.$$

Topological properties of the Poincaré map.

We use the following norm to compute the distances in each $\widehat{B}_k \subset B$, for $k \in I$:

$$\|V - W\| = \max_{i \in I} |V_i - W_i| \quad \forall V, W \in \widehat{B}_k \quad (10)$$

Assume the following generic hypothesis in the parameters of the system:

$$H_{ij} \neq \theta \quad \forall i \neq j \in I$$

Define the *expansivity constant*:

$$\alpha = \frac{\min_{i \neq j} |\theta - H_{ij}|}{4} > 0 \quad (11)$$

Lemma 2. *If $V \in \partial\mathcal{P}$ then the Poincaré map ρ is discontinuous in V and the discontinuity jumps in V are larger than 3α . Precisely, there exists a sequence of points $U^m \rightarrow V$ such that $\|\rho(U^m) - \rho(V)\| > 3\alpha$*

Proof: If $V \in \partial\mathcal{P}$ then $V \in B_J$ for some $J \in \wp(I)$, $\#J \geq 2$. There exist $i \neq j$ in J and therefore $B_J \subset \partial B_{\{i\}}$. By the definition of topological frontier, there exists a sequence of points $U^m \rightarrow V$ such that $U^m \in B_{\{i\}}$.

$$\rho_j(U^m) = \Phi_j^{\bar{t}(U^m)}(U^m) - H_{ij} \xrightarrow{m \rightarrow +\infty} \Phi_j^{\bar{t}(V)}(V) - H_{ij}$$

The last limit is computed recalling that the solution Φ^t of the differential equation (1) depends continuously on the initial state V , and the ISI time $\bar{t}(V)$, defined as the minimum in equality (3) of the solutions of the implicit equations (4), also depends continuously on V , due to the Implicit Function Theorem.

But $j \in J$, $V \in B_J \Rightarrow j \in J(V)$, or in other terms, V is an initial state for which the neuron j arrives to the threshold level. Then $\Phi_j^{\bar{t}(V)}(V) = \theta$, and by equalities (9), $j \in J(V) \Rightarrow \rho_j(V) = 0$. Then:

$$\begin{aligned} \lim \rho_j(U^m) &= \theta - H_{ij}, \quad \rho_j(V) = 0 \\ |\lim \rho_j(U^m) - \rho_j(V)| &= |\theta - H_{ij}| \geq 4\alpha > 3\alpha \end{aligned}$$

By the definition of limit, for all m large enough we obtain $\|\rho_j(U^m) - \rho_j(V)\| > 3\alpha$. \square

The Lemma 2 explains why the discontinuities of ρ act as saddle type orbits in continuous systems, to produce expansion and chaos. The points of $\partial\mathcal{P}$ act on their neighborhoods with an infinite rate of expansion in some directions, given by instantaneous expansivity larger than an uniform number $3\alpha > 0$. But they also have directions of contraction, as they are in the frontier of the open pieces $B_{\{i\}}$ of continuity of ρ , where the following result holds:

Theorem 3. *The Poincaré map $\rho : B \mapsto B$ is uniformly contractive in each of its continuity pieces $B_{\{i\}} \cap \rho(B)$. Precisely, there exists $0 < \lambda < 1$ and a distance $dist_c$ such that, if $V, U \in B_{\{i\}} \cap \rho(B)$ for some $i \in I$, then*

$$dist_c(V, U) \leq \lambda dist_c(V, U)$$

Besides, there exists $K > 0$ such that for all $V, U \in B$:

$$\frac{1}{K} \|V - U\| \leq dist_c(V, U) \leq \|V - U\|$$

We note that the distance dist_c is not necessarily the norm $\|\cdot\|$ defined in (10). Nevertheless, they are equivalent in \mathbb{R}^{n-1} . For simplicity we will omit the not zero constant factors $1/K$ and K in the forward computations along this work, and use $\|\cdot\|$ as if it were dist_c . This is not a restriction because, as a consequence of this Theorem, for any norm there exists an iterate p_0 such that ρ^{p_0} is contractive in its continuities pieces: $\|\rho^p(V) - \rho^p(U)\| \leq K \text{dist}_c(\rho^p(V), \rho^p(U)) \leq K \lambda^p \text{dist}_c(V, U) \leq K^2 \lambda^p \|V - U\| \leq (1/2) \|V - U\|$ for all p large enough.

Proof: Define $0 < \epsilon_0 = \min_{h \neq j} H_{h,j}$. Apply equality (9) and use, at the end of the following formulae, the Lagrange Mean Value Theorem to compute the difference of $\Phi_i^t(V)$, for different values of the time t , as the increment Δt multiplied by the derivative in an intermediate time T^m :

$$\begin{aligned}
V \in \rho(B_{\{k\}}) &\Rightarrow V_k = 0, \quad V_j \leq \theta - \epsilon_0 \quad \forall j \in I \\
V \in \widehat{B}_k &= \{V \in B : V_k = 0\} \\
V \in B_{\{i\}} &\Rightarrow \phi_i^{\bar{t}(V)}(V) = \theta, \quad \phi_i^0(V) = V_i \\
\epsilon_0 \leq \theta - V_i &= \bar{t}(V) d\Phi_i^t(V)/dt|_{t=T^m} \\
\epsilon_0 &\leq \bar{t}(V) F_i(\Phi^{T^m}(V)) \\
\Rightarrow \bar{t}(V) &\geq \epsilon_0 / \max_{1 \leq i \leq n} F_i(V) = t_0 > 0
\end{aligned} \tag{12}$$

By the Tubular Flux Theorem there exists a \mathcal{C}^1 bounded diffeomorphism which is a spatial change of variables $\xi : V \mapsto \widehat{V}$ from $Q \in \mathbb{R}^n$ onto $\widehat{Q} \in \mathbb{R}^n$, such that $\xi|_B = id$ and the solutions of the differential equation (1) in Q verify $d\widehat{V}/dt = \vec{a}$ in \widehat{Q} , where $\vec{a} \in \mathbb{R}^n$ is a constant vector with positive components. It verifies: $\xi(\phi^t(V)) = \xi(V) + \vec{a} \cdot t$, $d\xi F(V) = \vec{a} \quad \forall V \in Q$.

Define in R^n the orthogonal projection π onto the $(n-1)$ -dimensional subspace $a_1 V_1 + a_2 V_2 + \dots + a_n V_n = 0$. The flux of the differential equation (1), after the change ξ of variables in the space, is orthogonal to that subspace, and is transversal to \widehat{B}_k . Consider any real function g :

$$\begin{aligned}
&\forall \widehat{V}, \widehat{V} + d\widehat{V} \in \mathbb{R}^n : \\
&\pi(d\widehat{V}) = \pi(d\widehat{V} + \vec{a}g(\widehat{V}, d\widehat{V})). \\
&\forall V, V + dV, \quad U \in \widehat{B}_k, \text{ define} \\
&\text{dist}_c(V, V + dV) = \|\pi(d\xi_V dV)\| \\
\text{dist}_c(V, U) &= \int_0^1 \|\pi(d\xi_{V+t(U-V)} \cdot (U - V))\| dt
\end{aligned} \tag{13}$$

As ξ is a \mathcal{C}^1 diffeomorphism, its derivative and the derivatives of its inverse, are bounded in the compact set B , and so the distance dist_c defined above verifies the last thesis of this Theorem. It is left to prove that ρ is contractive with this distance in $B_{\{i\}} \cap \widehat{B}_k$.

Let us apply ρ to $V, V + dV \in B_{\{i\}} \cap \widehat{B}_k$. We use the equality (9) with $J(V) = \{i\}$, in which for convenience we agree to define $H_{i,i} = \theta$. We shall use the derivation formula of the flux of the differential equation respect to its initial state:

$$\begin{aligned}
&\partial \Phi_j^t(V) / \partial V_j = \exp \left(\int_0^t (\partial F_j / \partial V_j) (\Phi^s(V)) ds \right) \\
\text{Define: } -\gamma &= \max_{1 \leq j} \max_{V \in Q} \partial F_j(V) / \partial V_j < 0
\end{aligned}$$

The formula (9) and the inequality (12) lead to:

$$\begin{aligned}
\rho(V) - \rho(V + dV) &= d\rho_V \cdot dV = \\
&= [(\partial\rho_j(V)/\partial V_j)dV_j + (\partial\rho_j(V)/\partial V_i)dV_i]_{1 \leq j \leq n} \\
\partial\rho_j(V)/\partial V_j &= (\partial\Phi_j^t(V)(V)/\partial V_j)\Big|_{t=\bar{t}(V)} = \\
&= \exp\left(\int_0^{\bar{t}(V)} (\partial F_j/\partial V_j)(\Phi_j^s(V)) ds\right) \leq e^{-\gamma t_0} \\
\partial\rho_j(V)/\partial V_i &= \\
&= (d\Phi_j^t(V_j)(V)/dt)\Big|_{t=\bar{t}(V)} \cdot (d\bar{t}(V_i)/dV_i) = \\
&= g(V) \cdot F_j(\Phi^{\bar{t}(V)}(V))
\end{aligned} \tag{14}$$

where $g(V) = d\bar{t}(V_i)/dV_i$ is the real function obtained deriving respect to V_i the implicit equation $\theta = \Phi_i^{\bar{t}(V_i)}(V_i)$. Call \vec{e}_j to the j -th. vector of the canonic base in \mathbb{R}^n and join all the results above:

$$\begin{aligned}
\pi \cdot d\xi_{\rho(V)}(\rho(V) - \rho(V + dV)) &= \pi \cdot d\xi_{\rho(V)}d\rho_V \cdot dV = \\
&= \pi \cdot d\xi_{\rho(V)}\left(\sum_{j=1}^n (\partial\rho_j(V)/\partial V_j) \cdot dV_j \vec{e}_j\right) + \\
&\quad + \pi \cdot d\xi_{\rho_j(V)}(g(V)F(\phi^{\bar{t}(V)}(V))) = \\
&= \pi \cdot d\xi_{\rho_j(V)}\left(\sum_{j=1}^n (\partial\rho_j(V)/\partial V_j) \cdot dV_j \vec{e}_j\right) + \\
&\quad + g(V)\pi \cdot d\xi_{\rho(V)}F(\phi^{\bar{t}(V)}(V)) = \\
&= \pi \cdot d\xi_{\rho(V)}\left(\sum_{j=1}^n (\partial\rho_j(V)/\partial V_j) \cdot dV_j \vec{e}_j\right) + \\
&\quad + \pi(g(V)\vec{a}) = \\
&= \pi \cdot d\xi_{\rho(V)}\left(\sum_{j=1}^n (\partial\rho_j(V)/\partial V_j) \cdot dV_j \vec{e}_j\right)
\end{aligned} \tag{15}$$

Now we define the number $0 < \lambda = e^{-\gamma t_0} < 1$ and observe from the computations above that:

$$0 < \partial\rho(V)/\partial V_j \leq e^{-\gamma t_0} = \lambda < 1$$

Applying the definition of the differential distance dist_c in (13), and the equality (15), we obtain:

$$\begin{aligned}
\text{dist}_c(\rho(V), \rho(V + dV)) &= \|\pi(d\xi_{\rho(V)}d\rho_V \cdot dV)\| \leq \\
&\leq \lambda \|\pi d\xi \cdot dV\| = \lambda \text{dist}_c(V, V + dV) = \lambda \|\pi(d\xi_V dV)\|
\end{aligned}$$

Integrating by formula (13) we conclude:

$$\text{dist}_c(\rho(V), \rho(U)) \leq \lambda \text{dist}_c(V, U) \quad \square$$

Measure properties of the Poincaré map.

Let m be the $(n - 1)$ -dimensional Lebesgue probability measure in the Poincaré section B . Let us prove that

$$m(\partial\mathcal{P}) = m(\bigcup_{\#J \geq 2} B_J) = 0.$$

In fact, each B_J , if $\#J \geq 2$, is the finite union of C_2^n $n - (\#J)$ -dimensional manifolds in B , obtained when, for two or more different values of $i \in I$, the respective solutions $t_i(V)$ of the implicit equations (4), are equal.

Our aim is to study the attractors. The weakest condition required to a set $A \subset B$ to be an attractor is that its basin of attraction has positive Lebesgue measure. So, we may take out the points of the measure zero set $\partial\mathcal{P}$.

The set B' of the points with infinite itinerary.

Define

$$\begin{aligned} B' &= \{V \in B : \rho^j(V) \notin \partial\mathcal{P} \ \forall j \geq 0\} \subset B \setminus \partial\mathcal{P} \\ B_{\{i\}} \cap B_{\{j\}} &= \emptyset \text{ if } i \neq j \text{ and } \bigcup_{i \in I} B_{\{i\}} = B \setminus \partial\mathcal{P} \Rightarrow \\ B' &= \bigcap_{p=0}^{+\infty} \rho^{-p}(B' \setminus \partial\mathcal{P}), \quad B \setminus B' = \bigcup_{p=0}^{+\infty} \rho^{-p}(\partial\mathcal{P}). \end{aligned}$$

The following assertion characterizes the set B' :

$$V \in B' \Leftrightarrow \exists \text{ a unique sequence } \{i_p\}_{p \geq 0} \text{ such that } \rho^p(V) \in B_{\{i_p\}} \ \forall j \geq 0.$$

This sequence is called *the itinerary of V* , and it is the infinite sequence of neurons that will spike (in different times), in the order they reach the threshold potential, from the initial state V .

ρ is continuous in each $B_{\{i\}}$. Therefore, for all $p \geq 0$ and for all $V \in B'$, the iterate map ρ^p is continuous in V . As B' is the numerable intersection of open and dense sets, it is dense.

Theorem 4. *The set B' with infinite itinerary has full Lebesgue measure in B , i.e: $m(B \setminus B') = 0$.*

Proof: To prove that $m(\rho^{-1}(\partial\mathcal{P})) = 0$, once it is known that $m(\partial\mathcal{P}) = 0$, use the same argument as to prove by induction that for all $p \geq 1$: $m(\rho^{-p}(\partial\mathcal{P})) = 0$, once it is known for $p - 1$.

In fact, we apply Liouville Theorem to compute the following Lebesgue integral:

$$\begin{aligned} 0 = m(\partial\mathcal{P}) &\geq m(\rho(\rho^{-1}(\partial\mathcal{P}))) = \\ &= \int_{V \in \rho^{-1}(\partial\mathcal{P})} |\det(D\rho)(V)| \, dm(V) \end{aligned}$$

and conclude that $m((\rho^{-1}(\partial\mathcal{P})) \cap \{V \in B; |\det(D\rho)| > 0\}) = 0$. To prove that $m(\rho^{-1}(\partial\mathcal{P})) = 0$ it is enough to prove that the Jacobian $\det(D\rho) \neq 0$ a.e. in B .

To apply the Liouville Theorem we shall first prove that ρ , given by formulas (9), is differentiable a.e. in the integration set. The first technical problem arises because ρ is not differentiable in all the points: it is neither in the points of discontinuity $\partial\mathcal{P}$ nor in the points of $\bigcup_{h \neq k \in I} (\widehat{B}_h \cap \widehat{B}_k)$. In fact, in these last set the Poincaré section itself fails to be a local differentiable manifold. The set of those exceptional points has zero Lebesgue measure because they are contained in $(n - 2)$ -dimensional subspaces.

The second technical difficulty is to check that $\det D\rho \neq 0$ a.e. The derivative of the Poincaré map can be computed directly from formulas (9), in each continuity piece $B_{\{i\}}$ intersected with each $\widehat{B}_k \setminus \bigcup_{h \neq k} \widehat{B}_h$. In those sets B is a local differentiable manifold. Computations in (14) lead to

$\det D\rho(V) = \left[\prod_{j \neq i, j \neq k} (\partial\rho_j(V)/\partial V_j) \right] \cdot (\partial\rho_k(V)/\partial V_i)$ All these factors are computed in equalities (14) and are not zero due to hypothesis (2), so $\det D\rho \neq 0$ a.e. \square

3. CHAOTIC AND STABLE SETS.

We will divide the Poincaré section B in two complementary sets S , formed by stable orbits, and $C = B \setminus S$, the chaotic ones, according to Definition 5. The set S is formed by stable future orbits under any uniform sufficiently small perturbation that can be added at any step of the iteration of ρ . In the set $C = B \setminus S$ there are arbitrarily small perturbations that, if added in some step of the iteration of ρ , drastically change the future orbits and their asymptotic behavior. To have a criteria of chaos, what we call drastic changes in the phase state, we consider the expansivity constant $\alpha > 0$ defined in (11).

Definition 5. Stable and chaotic sets. $V \in B$ is *stable* if there exists $\delta > 0$ such that $\forall p \geq 0$, if $\|\rho^p(V) - W\| \leq \delta$ then $\|\rho^k(\rho^p(V)) - \rho^k(W)\| \leq \alpha \forall k \geq 1$.

$V \in B$ is *chaotic* if it is not stable. The opposite of the definition of stable point holds as follows: $V \in B$ is *chaotic* if and only if for all $\delta > 0$ there exists $p \geq 0$, and there exists $W \in B$ such that $\|\rho^p(V) - W\| \leq \delta$ and $\|\rho^k(\rho^p(V)) - \rho^k(W)\| > \alpha$ for some $k \geq 1$.

S is the set of all stable points and $C = B \setminus S$ is the set of all chaotic points.

It is immediate from the definitions above that S is forward invariant: $\rho(S) \subset S$, and thus, its complement C is backward invariant: $\rho^{-1}(C) \subset C$.

Given $\delta > 0$ fixed, we define the uniform stable set $S_\delta \subset S$ as the set of $V \in B$ such that $\forall p \geq 0$, if $\|\rho^p(V) - W\| \leq \delta$ (with δ previously fixed), then $\|\rho^k(\rho^p(V)) - \rho^k(W)\| \leq \alpha \forall k \geq 1$. From the definitions above observe that $\rho(S_\delta) \subset S_\delta$ and that any point in S is in S_δ for some $\delta > 0$. Therefore, to study the dynamics of the stable points, it is enough to study the dynamics in the sets S_δ .

Theorem 6. The set C of chaotic points is never empty. Precisely: $C \supset B \setminus B' \supset \partial\mathcal{P} \neq \emptyset$, where B' is the set of points in the Poincaré section B with infinite itinerary, and \mathcal{P} is the set of discontinuities of ρ .

Even more, if $V \in \partial\mathcal{P}$ then for all $\delta > 0$ there exists $W \in B$ such that $\|V - W\| < \delta$, $\|\rho(V) - \rho(W)\| > 3\alpha$, where α is the expansivity constant of the system.

Proof: From the note at the end of Remark 1 and from the definition of the set B' of the points with infinite itinerary, we get $\emptyset \neq \partial\mathcal{P} \subset B \setminus B'$.

First, let $V \in \partial\mathcal{P}$. We shall prove that $V \in C$. From Lemma 2 there exists sequence of points $U^m \rightarrow V$ such that $\lim \|\rho(U^m) - \rho(V)\| > 3\alpha$. Given $\delta > 0$ for all $m \geq 1$ large enough the points $U^m \rightarrow V$ verify $\|U^m - V\| < \delta$. Therefore, taking $W = U^m$ the point V verifies the definition of chaotic point with $p = 0$ and $k = 1$.

Second, take $V \in B \setminus B' = \bigcup_{p \geq 0} \rho^{-p}(\partial\mathcal{P})$. We shall prove that $V \in C$. We know that for some $p \geq 0$ $\rho^p(V) \in \partial\mathcal{P}$. In the first step we proved $\rho^p(V) \in C$, so $V \in \rho^{-p}(C) \subset C$, because C is backward invariant. \square

Classification of systems.

► a.e.stable systems: $m(S) = 1$ and $m(C) = 0$. In Theorem 8 we prove that the a.e the limit set is composed only by periodic sinks (limit cycles). Nevertheless the set C of chaotic points is not empty as proved in Theorem 6.

► Chaotic systems: $m(C) = 1$ and $m(S) = 0$. By definition the limit set A of C is a *chaotic attractor*. In Catsigeras & al. [2008] it is proved that A is a Cantor set attractor.

► Partially chaotic and partially periodic systems: $m(C) > 0$ and $m(S) > 0$. We will show that the points in S are attracted to limit cycles.

Remark: We did not find \mathcal{C}^1 examples in our model of systems for which $m(C) > 0$ although due to Theorem 6 the set C of chaotic points is never empty. Nevertheless it is possible (but not immediate) to construct piecewise continuous \mathcal{C}^0 maps ρ in a $n - 1 \geq 2$ compact ball B , that are uniformly contractive in each of their continuity pieces and such that $C = B$, $S = \emptyset$.

Some known results.

Suppose allowed any small perturbation of the system structure, such that, instead of having the differential vectorial equation (1), we have $dV/dt = G(V)$ with $\|G - V\|_{\mathcal{C}^1} < \epsilon$ for some $\epsilon > 0$ sufficiently small. By continuity in the \mathcal{C}^1 topology of the functional space, the first two assumptions (2) are also verified by G and the third one transforms in $\partial G_i / \partial G_j \approx 0 \quad \forall j \neq i$.

Besides, instead of restricting to a constant matrix $H_0 = [H_{ij}]_{i,j \in I}$ to describe the synaptic interactions, allow a matrix $H(V)$ of negative numbers $H(\cdot)$ to be a \mathcal{C}^1 function such that $\|H - H_0\|_{\mathcal{C}^1} < \epsilon$.

The new Poincaré map ρ will move respect to the old one. The new set of discontinuities $\partial\mathcal{P}$ of the Poincaré map will be (with the Hausdorff distance) near the old one, because its points are those defined by implicit functions, whose equations have \mathcal{C}^1 dependence on $G, H \in \text{cal}\mathcal{C}^1$. So they are near the old ones if $\|G - F\|_{\mathcal{C}^1}$ and $\|H - H_0\|_{\mathcal{C}^1}$ are small enough. We conclude that, if $\epsilon > 0$ is sufficiently small, the new Poincaré map will be \mathcal{C}^1 near the old one, in each of its continuity pieces.

In this scenario, the following results are known, leading to the genericity of the property $m(S) = 1, m(C) = 0$.

► If provided the additional open hypothesis of separation (pairwise disjointness of the closure of the images of the different pieces of continuity for some iterate of ρ), then Catsigeras & al.[2008] proved that generically (open and dense in the topology defined above), there exists a finite number of limit cycles that attracts all the orbits of B' . We conclude that $S = B'$, and thus $m(S) = 1$.

► If the contractive map ρ is affine in each of its continuity pieces, then Cessac [2008] proved that for generic values of the real parameters, not only in the topological sense (open and dense family of systems), but also in the Lebesgue measure sense (in the space of finite number of the real affinity parameters), there exists a finite number of limit cycles, that attracts all the orbits of B' . We conclude that $S = B'$ and $m(S) = 1$.

► If the contractive map ρ is not affine, but the matrix of interactions H is constant, Cessac & Viéville [2008] proved that either there exists limit cycles attracting the points of B' or the dynamics has positive entropy.

Definition 7. Omega limit and limit cycles.

For any $V \in B$, its *omega-limit* $\omega(V) \in B$ is the set of limit points of the future orbit of V . Precisely:

$$\omega(V) = \{W \in B : \exists p_j \rightarrow +\infty \text{ such that } \rho^{p_j}(V) \rightarrow W\}$$

As the phase space B is compact, the omega limit set of any point is not empty. The Poincaré map ρ is not continuous, so $\omega(V)$ is not necessarily forward invariant. Nevertheless the omega limit set is the same for all the points in the same orbit, as it is easy to check.

A set $L \in B$ is a *limit cycle* (also called a *periodic sink*) if it is a single periodic orbit and its *basin attraction* $B(L)$ contains an open neighborhood of L , being:

$$L = \{V, \rho(V), \dots, \rho^{r-1}(V)\}, \quad \rho^r(V) = V \quad r \geq 1$$

$$B(L) = \{W \in B : \omega(W) = L\}$$

Theorem 8. *If for some $\delta > 0$ the set S_δ of uniform stable points is not empty then, there exists a finite number N of limit cycles L_1, \dots, L_N such that the union $\bigcup_{k=1}^N B(L_k)$ of their basins of attractions includes S_δ .*

Remark: Let us show the main consequence of this theorem: if the set S of stable points has positive measure, it is not empty and, because all its points are included in uniformly stable sets S_δ , they all are attracted to limit cycles. By definition of limit cycles, the basins of attraction of different limit cycles are pairwise disjoint containing open sets. The topology in B has numerable basis, so we conclude that *there are at most a numerable quantity of limit cycles attracting all the orbits of S* . If besides $m(S) = 1$, then *a.e. point in B is attracted to a limit cycle*.

We will prove Theorem 8 at the end of the next Section.

4. THE ATOMIZATION OF THE SPACE.

Let us consider a system for which the stable set S is not empty. All the results in this Section will hold under this hypothesis.

As observed at the end of Definition 5: $S \neq \emptyset \Rightarrow S_\delta \neq \emptyset$ for some $\delta > 0$, and thus also for all sufficiently small $\delta > 0$. Fix $0 < \delta < \alpha$ such that $S_\delta \neq \emptyset$, where α is the expansivity constant. We recall that $S_\delta \subset B'$ from Theorem 6, and $\rho(S_\delta) \subset S_\delta$. Therefore the image $\rho^p(S_\delta)$ is disjoint with $\partial\mathcal{P}$ for all $p \geq 0$. So it is partitioned in n disjoint pieces (may be some of them are empty) when intersecting S_δ with $\{B_{\{i\}}\}_{i \in I}$. In each of this pieces ρ is continuous, and $\rho(S_\delta) = \bigcup_{i \in I} \rho(S_\delta) \cap B_{\{i\}}$.

Definition 9. Atoms of generation p. Given $i_1 \in I$ we call *atom of generation 1* to the following set:

$$A_{i_1} = \rho(S_\delta \cap B_{\{i_1\}})$$

There are at least one and at most $n = \#I$ non empty atoms of generation 1. Besides

$$\rho(S_\delta) = \bigcup_{i_1 \in I} A_{i_1}.$$

Given $(i_1, i_2) \in I \times I$ we call *atom of generation 2* to the following set: $A_{i_1, i_2} = \rho(A_{i_1} \cap B_{\{i_2\}})$.

There are at least one and at most $n^2 = (\#I)^2$ non empty atoms of generation 2. Besides $\rho^2(S_\delta) = \bigcup_{(i_1, i_2) \in I^2} A_{i_1, i_2}$.

By induction, if defined the atoms $\{A_{i_1, i_2, \dots, i_p}\}_{(i_1, \dots, i_p) \in I^p}$ of generation p , we define the *atoms of generation $p+1$* :

$$A_{i_1, i_2, \dots, i_p, i_{p+1}} = \rho(A_{i_1, i_2, \dots, i_p} \cap B_{\{i_{p+1}\}})$$

$$\text{Then: } \rho^p(S_\delta) = \bigcup_{(i_1, i_2, \dots, i_p) \in I^p} A_{i_1, i_2, \dots, i_p}.$$

There are at least one and at most $n^p = (\#I)^p$ non empty atoms of generation p . Fix $p \geq 1$. Denote \mathcal{A}_p to the finite collection of atoms of generation p : $\mathcal{A}_p = \{A_{i_1, i_2, \dots, i_p}\}_{(i_1, \dots, i_p) \in I^p}$.

Remark: It is easy to check from the Definition above that for all $p \geq 1$, if $A \in \mathcal{A}_{p+1}$ then there exists $A' \in \mathcal{A}_p$ such that $A \subset A'$. Precisely $A = A_{i_1, i_2, \dots, i_{p+1}} \subset A_{i_2, \dots, i_{p+1}} = A'$

We define the *diameter d_p of the atomization* of generation p : $d_p = \max_{A \in \mathcal{A}_p} \text{diam}(A)$ where $\text{diam}(A) = \sup_{V, W} \|V - W\|$ denotes the diameter of the atom A and it is 0 if $A = \emptyset$. We observe the d_p is not the diameter of the union of the atoms, even if they intersect.

Lemma 10. *Let $0 < \lambda < 1$ the uniform contraction rate of Theorem 3. There exists a constant K such that for any $p \geq 1$ the diameter d_p of the atomization of generation p verifies:*

$$0 \leq d_p \leq K\lambda^{p-1}, \quad \lim_{p \rightarrow +\infty} d_p = 0.$$

Proof: Take $K = \text{diam}B$. Let us prove the thesis by induction on $p \geq 1$.

An atom $A \in \mathcal{A}_1$ of generation 1 is the image by ρ of a set in B . Then $\text{diam}(A) \leq K$. Suppose that all the atoms of generation p have diameter smaller than $K\lambda^{p-1}$. Let us prove the thesis for $p + 1$. An atom $A \in \mathcal{A}_p$, $p \geq 1$ is the image by ρ of some set in B , so $A \subset \rho(B)$. Take an atom $A' \in \mathcal{A}_{p+1}$. By definition: $A' = \rho(A \cap B_{\{i\}})$ for some $A \in \mathcal{A}_p$ and some $i \in I$.

As $A \cap B_{\{i\}} \subset \rho(B) \cap B_{\{i\}}$ due to Theorem 3 ρ is contractive there:

$$\begin{aligned} \text{diam}(A') &= \sup_{V^1, V^2 \in A'} \|V^1 - V^2\| = \\ &= \sup_{U^1, U^2 \in A \cap B_{\{i\}}} \|\rho(U^1) - \rho(U^2)\| \leq \\ &\leq \sup_{U^1, U^2 \in A \cap B_{\{i\}}} \lambda \|\rho(U^1) - \rho(U^2)\| = \\ &= \lambda \text{diam}(A \cap B_{\{i\}}) \leq \lambda \text{diam}(A) \leq \\ &\leq K \lambda (\lambda^{p-1}) = K \lambda^p. \quad \square \end{aligned}$$

Recall the distance $\text{dist}(V, A)$ of a point $V \in B$ to a non empty set $A \subset B$: $\text{dist}(V, A) = \sup_{W \in A} \|V - W\|$. Denote \overline{A} and ∂A to the closure and frontier respectively, of the set A . A basic classic result from Topology asserts that, in any compact connected metric space, the following properties hold: $\text{dist}(V, A) = \text{dist}(V, \overline{A}) = \text{dist}(V, \partial A) = \max_{W \in \partial A} \|V - W\| = \|V - U_0\|$ for some $U_0 \in \partial A$.

Lemma 11. *If $V \in S_\delta$ then $\text{dist}(V, \overline{\partial \mathcal{P}}) \geq \delta/2$.*

Proof: Due to the metric properties in compact connected spaces recalled above, it is enough to show that $\text{dist}(V, \partial \mathcal{P}) \geq \delta/2$. By contradiction suppose that $\inf_{U \in \partial \mathcal{P}} \|V - U\| < \delta/2$. Then there exists $U \in \partial \mathcal{P}$ such that:

$$\|V - U\| < \delta/2.$$

$V \in B_\delta$ and by definition of uniform stability, if $\|V - W\| < \delta$ then:

$$\|\rho^k(V) - \rho^k(W)\| \leq \alpha \quad \forall k \geq 1 \tag{16}$$

In particular $\|\rho(V) - \rho(U)\| \leq \alpha$.

On the other hand $U \in \partial \mathcal{P}$. Applying Theorem 6 there exists $W \in B$ such that

$$\|U - W\| < \delta/2, \quad \|\rho(U) - \rho(W)\| > 3\alpha.$$

Joining the inequalities above with the triangular property: $\|V - W\| \leq \|V - U\| + \|U - W\| < \delta$,

$$\begin{aligned} 3\alpha &< \|\rho(U) - \rho(W)\| \leq \\ &\leq \|\rho(U) - \rho(V)\| + \|\rho(V) - \rho(W)\| \leq \\ &\leq \alpha + \|\rho(V) - \rho(W)\|, \end{aligned}$$

$$\|V - W\| < \delta, \quad 2\alpha \leq \|\rho(V) - \rho(W)\|. \quad \text{These last two inequalities contradict (16).} \quad \square$$

Lemma 12. *If $V \in S_{\delta'} \cap B_{\{i\}}$, then for all $W \in B$ such that $\|W - V\| < \delta/2$ it is verified $W \in B_{\{i\}}$.*

Proof: Consider the compact set $K = B \setminus B_{\{i\}}$. Then:

$$U \in K \Leftrightarrow U \notin B_{\{i\}}.$$

We have $V \notin K$. Call $d = \text{dist}(V, K) = \inf_{U \notin B_{\{i\}}} \|V - U\|$. As recalled when defining the distance of a point to a set, there exists $U_0 \in \partial K = \partial U_{\{i\}} \subset \overline{\partial \mathcal{P}}$ such that

$$d = \|V - U_0\| = \text{dist}(V, \partial K) \geq \text{dist}(V, \overline{\partial \mathcal{P}}).$$

Due to Lemma 11 $\text{dist}(V, \overline{\partial \mathcal{P}}) \geq \delta/2$. In resume we have proved that: $d \geq \delta/2$, $\inf_{U \notin B_{\{i\}}} \|V - U\| \geq \delta/2$ Therefore, if $\|V - W\| < \delta/2$ then $W \in B_{\{i\}}$. \square

Definition. Indivisibility of the atoms. An atom $A \in \mathcal{A}_p$ is *indivisible* if its closure $\overline{A} \subset B_{\{i\}}$ for some $i \in I$. From the definition of atom, $\rho(A)$ is a unique atom of generation $p + 1$, i.e. A does not divide when applying ρ . Besides ρ is continuous in $B_{\{i\}}$ so, it is continuous in \overline{A} . Any atom $A \in \rho(B)$, so applying Theorem 3:

If an atom A is indivisible then ρ is continuous and uniformly contractive in \overline{A} .

Proof of Theorem 8:

By Lemma 10 there exists a generation $p \geq 1$ of atoms such that the diameter $d_p < \delta/2$. Applying Lemma 12 and recalling that the diameter of a set is the same diameter of its closure, each atom $A \in \mathcal{A}_p$ is indivisible. From the remark at the end of Definition 9 the atoms of generation $p + 1$ and later are contained in the atoms of generation p . Then all of them are also indivisible.

Fix some non empty atom $A = A^1 \in \mathcal{A}_p$. As it is indivisible: $\emptyset \neq \rho(A^1) \in \mathcal{A}_{p+1}$. But any atom of generation $p + 1$ is contained in an atom of generation p , so there exists $A^2 \in \mathcal{A}_p$ such that $\rho(A^1) \subset A^2$. From the indivisibility, ρ is continuous in A^1 , thus: $\rho(\overline{A^1}) \subset \overline{A^2}$. Applying the same argument to A_2 instead of A_1 there exists $A^3 \in \mathcal{A}_p$ such that $\rho(\overline{A^2}) \subset \overline{A^3}$. We deduce $\rho^2(\overline{A^1}) \subset \rho(\overline{A^2}) \subset \overline{A^3}$ where $A^1, A^2, A^3 \in \mathcal{A}_p$. The family \mathcal{A}_p is finite, so there is some first pair of integer numbers $k_0 \geq 1, r_0 \geq 1$ such that $A^{k_0} = A^{r_0+k_0} \in \mathcal{A}_p$. We conclude that some subfamily of \mathcal{A}_p is related in a finite chain:

$$\begin{aligned} \rho(\overline{A^1}) \subset \overline{A^2}, \dots, \rho(\overline{A^j}) \subset \overline{A^{j+1}}, \dots, \\ \dots \rho(\overline{A^{k_0}}) \subset \overline{A^{k_0+1}}, \dots, \\ \dots \rho(\overline{A^{k_0+r_0-1}}) \subset \overline{A^{k_0+r_0}} = \overline{A^{k_0}} \\ \rho^{r_0}(\overline{A^{k_0}}) \subset \overline{A^{k_0}} \end{aligned} \tag{17}$$

All these atoms are indivisible by construction, so for each $\overline{A^j}$, $j = 1, 2, \dots, \leq r_0 - 1$ in the finite chain (17), the Poincaré map $\rho : \overline{A^j} \mapsto \overline{A^{j+1}}$ is continuous and contractive, with a uniform contraction rate $0 < \lambda < 1$. In resume we have $\rho^{r_0} : \overline{A^{k_0}} \mapsto \overline{A^{k_0}}$ and

$$\|\rho^{r_0}(V) - \rho^{r_0}(W)\| \leq \lambda^{r_0} \|V - W\| \text{ for all } V, W \in \overline{A^{k_0}}.$$

The Brower Theorem of the Fixed Point, states that in any compact metric space M , given a uniformly contractive map f such that $f(M) \subset M$, there exists and is unique a point $x_1 \in M$ fixed by f : $f(x_1) = x_1$. Besides all orbits by future iterates of f have $\lim_{k \rightarrow +\infty} \{x_1\}$, i.e. $\lim_{k \rightarrow +\infty} f^k(x) = x_1$, $\forall x \in M$. We conclude that ρ^{r_0} in the compact metric space $\overline{A^{k_0}}$ has a fixed point V^1 such that:

$$\begin{aligned} V^1 \in \overline{A^{k_0}}, \quad \rho^{r_0}(V^1) = V^1, \\ \lim_{k \rightarrow +\infty} \rho^{k r_0}(V) = V_1 \quad \forall V \in \overline{A^{k_0}} \end{aligned}$$

Therefore V^1 is periodic by ρ of period r_0 , and its orbit $L_1 = \{V^1, \rho(V^1), \dots, \rho^{r_0-1}V^1\}$ is the omega limit of all the points in $\bigcup_{j=1}^{j=k_0+r_0} \bar{A}^j$, in particular of those in \bar{A}^1 .

From the definition of atom and from $\rho(S_\delta) \subset S_\delta$ each atom is contained in S_δ . By Lemma 12 and using that ρ is contractive in each of its continuity pieces intersected with $\rho(B)$, we deduce that all the points W such that $\text{dist}(\rho(W), \bigcup_{j=k_0+1}^{k_0+r_0} \bar{A}^j) < \delta/2$ have $\omega(W) = L_1$. Such points W form an open set $\mathcal{N} \supset \bigcup_{j=k_0}^{k_0+r_0-1} \bar{A}^j$. Besides $L_1 \subset \bigcup_{j=k_0}^{k_0+r_0-1} \bar{A}^j$. Then \mathcal{N} is a neighborhood of L_1 . Therefore the basin of attraction $B(L_1)$ of the periodic orbit L_1 contains a neighborhood of itself, verifying the definition of limit cycle.

In particular, $B(L_1) \supset \bar{A}^1 \in \mathcal{A}_p$. The construction above can be done choosing any first atom $A_1 \in \mathcal{A}_p$. We deduce that for each $A \in \mathcal{A}_p$ there exists a limit cycle $L \subset B'$ such that $\bar{A} \subset B(L)$.

Define $\mathcal{L} = \{L \subset B : L \text{ is a limit cycle, } A \subset B(L) \text{ for some } A \in \mathcal{A}_p\}$. It is a finite and not empty collection because the family \mathcal{A}_p of the atoms of generation p is finite and not empty. By construction the union of the basin of attractions of all the limit cycles in \mathcal{L} contains all the atoms of generation p . Therefore:

$\rho^p(B_\delta) = \bigcup_{A \in \mathcal{A}_p} A \subset \bigcup_{L \in \mathcal{L}} B(L)$. For all U the limit set $\omega(U) = \omega(\rho^p(U))$. Then $B_\delta \subset \bigcup_{L \in \mathcal{L}} B(L)$. \square

5. CONCLUSIONS

We described and analyzed a general mathematical model of a network of n inhibitory pacemaker neurons interacting by synapsis without delay. The size of the network must be finite but as large as wanted. We found discontinuities in the dynamical system due to the synaptic coupling and proved that they imply the existence of chaotic orbits. The Poincaré return map ρ to the $(n-1)$ -dimensional section B in the phase space, results from considering the state of the system immediately after each spike. We proved topological and measure properties of ρ as mathematical tools to obtain the dynamical results. We classified the systems according to the measure of the chaotic set and proved that, even being this set never empty, if it has not full measure there exist stable points and thus, limit cycles. It is unknown if there exist \mathcal{C}^1 systems of this model exhibiting a set of chaotic orbits with positive measure, although under some other additional hypothesis, there are known results about the genericity of the systems exhibiting only limit cycles.

Acknowledgments

We thank the project PDT 54/001 of Clemente Estable fund (Uruguay), the University of Valparaíso (Chile) and the University of Marburg (Germany) for partial support, Profs. Pierre Guiraud, Hans Braun and Ruben Budelli for their suggestions, and MEDYFINOL organizing & scientific comitee for its invitation.

REFERENCES

- Abbott, L.F. & Vreeswijk. C [1993] “ Assynchronous states in neural networks of pulse-coupled oscillators.” *Phys. Rev. E* **48** 1483-1490
- Bessloff, P. & Coombes, S. [2000] “ Dynamics of Strongly Coupled Spiking Neurons.” *Neural Computation* **12** 91-129.
- Budelli, R. , Catsigeras, E. , Rovella, A. & Gómez, L. [1996] “ Dynamical behavior of pacemaker neurons networks.” *Proc. of the Second Congress of Nonlinear Analysts, Elsevier Science.*
- Budelli, R., Torres, J., Catsigeras, E. , & Enrich, H. [1991] “ Two neurons networks I: Integrate and fire pacemakers models.” *Biol. Cybern.* **66**, 95-110

Catsigeras, E. & Budelli, R. [1992] “Limit cycles of a bineuronal network model.” *Physica D* **56**, 235-252

Catsigeras, E., Rovella, A. & Budelli, R. [2008] “Contractive piecewise continuous maps modeling networks of inhibitory neurons.” *ArXiv [q-bio NC]* **0805.2695v1**

Cessac, B. [2008] “A discrete time neural network model with spiking.” *J. Math. Biol.* **54**, 311-345

Cessac, B. & Viéville, T. [2008] “On Dynamics of integrate and fire neural networks with conductance based synapses.” *ArXiv [phys-bio-ph]* **0709.4370v3**

Mirollo, R.E. & Strogatz, S.H. [1990] “Synchronisation of pulse coupled biological oscillators” *SIAM, J. Appl. Math.* **50** 1645-1662

Postnova, S., Wollweber, B., Voigt, K., Braun, H. A. [2007] “Neural Impulse Pattern in Bidirectionally Coupled Model of Neurons of Different Dynamics.” *Biosystems* **89** 135-142