

Weak Pseudo-Physical Measures and Pesin's Entropy Formula for Anosov C^1 -diffeomorphisms.

Eleonora Catsigeras, Marcelo Cerminara, and Heber Enrich

Dedicated to the Memory of Prof. Nikolai Chernov

ABSTRACT. We consider C^1 Anosov diffeomorphisms on a compact Riemannian manifold. We define the weak pseudo-physical measures, which include the physical measures when these latter exist. We prove that ergodic weak pseudo-physical measures do exist, and that the set of invariant probability measures that satisfy Pesin's Entropy Formula is the weak*-closed convex hull of the ergodic weak pseudo-physical measures. In brief, we give in the C^1 -scenario of uniform hyperbolicity, a characterization of Pesin's Entropy Formula in terms of physical-like properties.

1. Introduction

The purpose of this paper is to give, in the C^1 -scenario of uniform hyperbolicity, a characterization of those invariant measures that satisfy Pesin's Entropy Formula in terms of their physical-like properties. Our main result works, for C^1 Anosov diffeomorphisms, as Ledrappier-Young characterization [LY] of the measures μ that satisfy Pesin's Entropy Formula (which holds in the C^2 context but not in the general C^1 context), by substituting the property of absolute continuity of the unstable conditional decomposition of μ , by the weak pseudo-physical property of its ergodic decomposition.

Pesin Theory [P1, P2] gives relevant tools and results of the modern differentiable ergodic theory. It works for C^2 (or at least C^1 plus Hölder) dynamical systems. For instance, for C^2 hyperbolic systems, Pesin's Entropy Formula computes exactly the metric entropy of a diffeomorphism in terms of the mean value

1991 *Mathematics Subject Classification.* Primary 37A35, 37D20; Secondary 37D35; 37A05.

Key words and phrases. Anosov diffeomorphisms; Pesin's Entropy Formula; Physical Measures; SRB Measures.

E.C. thanks the Department of Mathematics of the University of Alabama at Birmingham, Prof. Alexander Blokh, and the scientific and organization committees of *The Dynamical Systems, Ergodic Theory and Probability Conference Dedicated to the Memory of Nikolai Chernov*. E.C. is partially supported by Project "PhySeCo" of the program MathAmSud of France, Chile and Uruguay (through A.N.I.I. Uruguay).

©0000 (copyright holder)

of the sum of its positive Lyapunov exponents. In the C^2 scenario, Pesin's Entropy Formula holds if and only if the invariant measure has absolutely continuous conditional decomposition along the unstable manifolds. [LY].

Through the properties of absolute continuity of invariant measures, and mainly through the absolute continuity of the holonomy along the invariant foliations, Pesin Theory gives the tools to construct physically significant invariant measures for C^1 -plus Hölder systems. Among these measures, the so called Sinai-Ruelle-Bowen (SRB) measures [Si, R1, B1], have particular relevance to describe the asymptotic statistics of Lebesgue-positive set of orbits, not only for C^1 plus Hölder uniform and non-uniform hyperbolic systems, but also for C^1 plus Hölder partially hyperbolic systems [PS, BDV]. Precisely, one of the most relevant properties of ergodic SRB measures for C^1 plus Hölder hyperbolic systems, is that they are *physical*; namely, their basins of statistical attraction have positive Lebesgue measure, even for Lebesgue non-preserving systems.

In particular, for transitive Anosov C^1 plus Hölder systems, the theorem of Pesin-Sinai (see for instance [PS]) states that there exists a unique physical measure: it is the unique invariant probability measure that satisfies Pesin's Entropy Formula, and so, the only one with absolutely continuous conditional measures along the unstable foliation. Besides, its basin of statistical attraction covers Lebesgue all the orbits. In other words, for a C^1 plus Hölder Anosov system, the definition of physical measure, SRB measure, and measure that satisfies Pesin's Entropy Formula, are equivalent.

Nevertheless, in the C^1 -scenario, the above results do not work, because the theorems of Pesin Theory that ensure the absolute continuity of unstable conditional measures, and of the holonomies along invariant foliations, fail. Even the existence of the unstable manifolds, along which one could construct the conditional unstable measures, fails in the C^1 context [Pu]. In the particular case of C^1 -Anosov diffeomorphisms, invariant C^0 foliations with C^1 leaves do exist (see for instance [HPS]), but the holonomies along the invariant foliations are not necessarily absolutely continuous [RoY].

As a consequence, for C^1 systems, if one defined SRB measures by the existence of their absolutely continuous unstable conditional measures, one would lack the hope to construct them. Nevertheless, one can still define SRB or SRB-like measures, if one forgets for a while the properties of absolute continuity, and focus the attention of the properties of statistical attraction. In other words, one can try to look directly at their physical properties, dodging the lack of conditional absolute continuity. For that reason, in the C^1 -scenario, we look for the Lebesgue abundance of points in their statistical basins (or, more precisely, in the ϵ -approach of their statistical basins). This search was used in [CE] to construct the SRB-like or pseudo-physical measures for C^0 dynamical systems on a compact Riemannian manifold. The notion of SRB-like or pseudo-physical measures, even in a non differentiable context, translate to the space of probability measures the concept of statistical attraction defined by Ilyashenko [GI1, II] in the ambient manifold.

In [Qiu] it is proved that C^1 *generically*, transitive and uniformly hyperbolic systems do have a unique measure satisfying Pesin's Entropy Formula. Besides this measure is physical and its basin of statistical attraction covers Lebesgue almost all the orbits. In the general C^1 -scenario with non uniform hyperbolicity, Pesin's

Entropy Formula was first proved considering systems that preserve a smooth measure (we call a measure μ smooth if $\mu \ll \text{Leb}$, where Leb is the Lebesgue measure). In fact, for C^1 generic diffeomorphisms that preserve a smooth measure μ , [Ta] proved that μ satisfies Pesin's Entropy Formula. Later, in [ST], this formula was also proved for any C^1 partially hyperbolic system that preserves a smooth measure.

If no smooth measure is preserved, in [CCE] is proved that the pseudo-physical or SRB-like measures still exist and satisfy Pesin's Entropy Formula, provided that the system is C^1 partially hyperbolic. Recently, and also for C^1 partially hyperbolic systems, [YC] derived a proof of Shub's Entropy Conjecture [Sh] from their method of construction of measures that satisfy Pesin's Entropy Formula.

In this paper we focus on C^1 Anosov systems to search for a converse of the result in [CCE]. Namely, our purpose is to characterize all the invariant measures that satisfy Pesin's Entropy Formula. First, we need to generalize the concept of pseudo-physical or SRB-like measure. So, we define the *weak pseudo physical* measures μ , by taking into account only the ϵ -approach of its basin of statistical attraction up to time n , which we denote by $A_{\epsilon,n}(\mu)$, and the exponential rate of the variation of the Lebesgue measure of $A_{\epsilon,n}(\mu)$ when $n \rightarrow +\infty$ (Definition 1.3). In Theorem 1 we study general properties of the weak pseudo physical measures, which do always exist. We prove that for any C^1 Anosov diffeomorphism, the weak pseudo-physical measures satisfy Pesin's Entropy Formula (Part A) of Theorem 2). Besides, we prove a converse result, to conclude that the set of invariant measures that satisfy Pesin's Entropy Formula is the closed convex hull of the weak pseudo-physical measures (Theorem 2).

So, Theorem 2 characterizes all the measures that satisfy Pesin's Entropy Formula in terms of the statistical properties that define the weak pseudo-physical notion. Nevertheless, as far as we know, no example is still known of a C^1 Anosov diffeomorphism for which weak pseudo-physical measures are not physical. In other words, there are not known examples of C^1 -Anosov systems such that an ergodic measure satisfies Pesin's Entropy Formula and is non physical.

The proof of Theorem 2 is based on the construction of local C^1 pseudo-unstable foliations, which approach the local C^0 unstable foliation, and allow us to apply a Fubini decomposition of the Lebesgue measure of the ϵ -basin $A_{\epsilon,n}(\mu)$, for any ergodic measure μ . The pseudo-unstable foliations are constructed via Hadamard graphs whose future iterates have bounded dispersion. This method was introduced by Mañé in [Mañ] to prove Pesin's Entropy Formula in the C^1 plus Hölder context. Much later, it was applied also to C^1 systems in [ST] and [CCE].

As said above, in Theorem 2 of this paper, we prove that the weak pseudo-physical condition for the ergodic components of an invariant measure, is necessary and sufficient to satisfy Pesin's Entropy Formula. The sufficient condition is just a corollary of the results in [CCE]. On the contrary, the proof of the necessary condition is new, although it is also strongly based on Mañé's method to construct, via Hadamard graphs, the C^1 -pseudo unstable foliations.

As a subproduct of the proof of Theorem 2, we also obtain an equality for any ergodic measure of a C^1 -Anosov diffeomorphism, even for measures that do not satisfy Pesin's Entropy Formula. This equality, which is stated in Theorem 3, considers the exponential rate

$$a(\mu) := \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon,n}(\mu))}{n},$$

according to which the Lebesgue measure of the ϵ -basin $A_{\epsilon,n}(\mu)$ of each ergodic measure μ varies with time n . Theorem 3 equals the exponential rate $a(\mu)$ with the difference

$$h_\mu(f) - \int \sum_i \chi_i^+ d\mu,$$

where $h_\mu(f)$ is the metric entropy and $\{\chi_i^+\}$ are the positive Lyapunov exponents. So, in the particular case of ergodic measures μ satisfying Pesin's Entropy Formula, the exponential rate $a(\mu)$ is null, and conversely.

1.1. Definitions and Statement of the Results. Let M be a compact, connected, Riemannian C^1 -manifold without boundary and let $f : M \mapsto M$ be continuous.

DEFINITION 1.1. (EMPIRIC PROBABILITY) For each $x \in M, n \in \mathbb{N}^+$, the *empiric probability* $\sigma_n(x)$ along the finite piece of the future orbit of x up to time n , is defined by

$$\sigma_n(x) := \frac{1}{n} \sum_{j=0}^{n-1} \delta_{f^j(x)},$$

where δ_y denotes the Dirac delta probability measure supported on the point $y \in M$.

We denote by \mathcal{M} the space of all the Borel probability measures on M , endowed with the weak* topology. We denote by $\mathcal{M}_f \subset \mathcal{M}$ the space of f -invariant Borel probability measures. It is well known that \mathcal{M} and \mathcal{M}_f are nonempty, weak* compact, metrizable, sequentially compact and convex topological spaces. We fix and choose a metric dist^* in \mathcal{M} that induces the weak* topology.

DEFINITION 1.2. (BASIN AND PSEUDO BASIN OF ATTRACTION OF A MEASURE.) Let $\mu \in \mathcal{M}$ and $\epsilon > 0$. We construct the following measurable sets in the manifold M :

$$(1.1) \quad B(\mu) := \left\{ x \in M : \lim_{n \rightarrow +\infty} \sigma_n(x) = \mu \right\};$$

$$(1.2) \quad A_\epsilon(\mu) := \left\{ x \in M : \liminf_{n \rightarrow +\infty} \text{dist}^*(\sigma_n(x), \mu) < \epsilon \right\};$$

$$(1.3) \quad A_{\epsilon,n}(\mu) := \left\{ x \in M : \text{dist}^*(\sigma_n(x), \mu) < \epsilon \right\}.$$

We call $B(\mu)$ the *basin of attraction* of μ . We call $A_\epsilon(\mu)$ the ϵ -*pseudo basin of attraction* of μ . We call $A_{\epsilon,n}(\mu)$ the ϵ -*pseudo basin of μ up to time n* .

In the sequel we denote by Leb the Lebesgue measure of M , renormalized to be a probability measure.

DEFINITION 1.3. (PHYSICAL, PSEUDO-PHYSICAL AND WEAK PSEUDO-PHYSICAL MEASURES)

Let $\mu \in \mathcal{M}$. We call μ *physical* if $\text{Leb}(B(\mu)) > 0$.

We call μ *pseudo-physical* if $\text{Leb}(A_\epsilon(\mu)) > 0$ for all $\epsilon > 0$.

We call μ *weak pseudo-physical* if

$$(1.4) \quad \limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb}(A_{\epsilon,n}(\mu)) = 0 \quad \forall \epsilon > 0.$$

We denote

$$\mathcal{P}_f := \{\mu \in \mathcal{M} : \mu \text{ is weak pseudo-physical}\}.$$

On the one hand, it is standard to check that for continuous mappings $f : M \mapsto M$ the physical measures, if they exist, are f -invariant. Also, pseudo-physical are f -invariant (see [CE]), page 153, and as proved in Theorem 1.3 of [CE], the set of pseudo-physical measures is never empty, weak* compact and independent of the chosen metric dist^* that induces the weak* topology of \mathcal{M} . Besides, it is immediate to check that physical measures, if they exist, are particular cases of the always existing pseudo-physical measures.

On the other hand, in this paper we will generalize the previous results that hold for pseudo-physical measures, by proving the following properties also for weak pseudo-physical measures:

THEOREM 1. *Let $f : M \mapsto M$ be a continuous map. Then:*

- A) *Weak pseudo-physical measures are f -invariant.*
- B) *Physical measures and pseudo-physical measures are particular cases of weak pseudo-physical measures.*
- C) *Weak pseudo-physical measures do always exist.*
- D) *The set \mathcal{P}_f of weak pseudo-physical measures does not depend on the choice of the metric dist^* that induces the weak* topology on \mathcal{M} .*
- E) *\mathcal{P}_f is weak*-compact, hence sequentially compact.*
- F) *$\lim_{n \rightarrow +\infty} \text{dist}^*(\sigma_n(x), \mathcal{P}_f) = 0$ for Lebesgue almost all $x \in M$.*
- G) *If the weak pseudo-physical measure μ is unique, then it is physical and its basin of attraction $B(\mu)$ covers Lebesgue a.e. $x \in M$.*

REMARK 1.4. Weak pseudo-physical measures are not necessarily ergodic (see Example 5.4 of [CE]).

Now, let $f \in \text{Diff}^1(M)$ be a C^1 diffeomorphism on M .

DEFINITION 1.5. (ANOSOV DIFFEOMORPHISMS) The diffeomorphism f is called *Anosov* if there exists a Riemannian metric of M and a splitting $TM = E \oplus F$ which is continuous and non trivial (i.e. $\dim(E), \dim(F) \neq 0$), and a constant $\lambda < 1$, such that

$$(1.5) \quad \|Df_x|_{E(x)}\|, \quad \|Df_x^{-1}|_{F(x)}\| \leq \lambda \quad \forall x \in M.$$

We call E and F the *stable and unstable subbundles* respectively. We call λ the (*uniform*) *hyperbolicity constant*.

REMARK 1.6. We observe that the condition of continuity of the unstable and stable subbundles is redundant in Definition 1.5. Besides, since the manifold is connected, from the continuity of F and E we deduce that they are uniformly transversal sub-bundles and $\dim(F)$ and $\dim E$ are constants.

From inequalities (1.5), for any Anosov diffeomorphism f and for any regular point $x \in M$, the minimum Lyapunov exponent along $F(x)$ is not smaller than $\log \lambda^{-1} > 0$, and the maximum Lyapunov exponent along E is not larger than $\log \lambda < 0$. Thus, for any regular point $x \in M$ all the Lyapunov exponents along $E(x)$ are strictly negative and bounded away from zero, and all the Lyapunov exponents along $F(x)$ are strictly positive and bounded away from zero.

DEFINITION 1.7. (PESIN'S ENTROPY FORMULA) Let $f \in \text{Diff}^1(M)$. Let $\mu \in \mathcal{M}_f$. We say that μ satisfies Pesin's Entropy Formula if

$$h_\mu(f) = \int \sum_{i=1}^{\dim(M)} \chi_i^+(x) d\mu,$$

where $h_\mu(f)$ is the metric entropy of f with respect to μ ; for μ -a.e. $x \in M$ the Lyapunov exponents of the orbit of x are denoted by

$$\chi_1(x) \geq \chi_2(x) \geq \dots \geq \chi_{\dim M}(x);$$

and $\chi_i^+(x) := \max\{\chi_i(x), 0\}$.

Recall that for any C^1 -Anosov diffeomorphism f , the set of measures that satisfy Pesin's Entropy Formula is nonempty (see for example Theorems 4.2.3 and 4.5.6 of [Ke]).

The main purpose of this paper is to prove the following result:

THEOREM 2. *For C^1 Anosov diffeomorphisms, the set of ergodic weak pseudo-physical measures is nonempty, and the set of invariant probability measures that satisfy Pesin's Entropy Formula is its closed convex hull.*

The following is an equivalent restatement of Theorem 2:

- A) *All the weak pseudo-physical measures satisfy Pesin's Entropy Formula.*
- B) *Any invariant probability measure μ satisfies Pesin's Entropy Formula if and only if its ergodic components μ_x are weak pseudo-physical μ -a.e. $x \in M$.*

From Theorem 2, we obtain the following consequence:

COROLLARY 1.8. *If $f \in \text{Diff}^1(M)$ is Anosov, then for Lebesgue-almost all $x \in M$ any convergent subsequence of the empirical probabilities $\sigma_n(x)$ converges to a measure that satisfies Pesin's Entropy Formula.*

PROOF. From Assertion F) of Theorem 1, for Lebesgue-almost all $x \in M$ any convergent subsequence of $\{\sigma_n(x)\}_{n \geq 1}$ converges to a weak pseudo-physical measure μ . Thus, applying part A) of Theorem 2 μ satisfies Pesin's Entropy Formula. \square

The arguments to prove Theorem 2 are based in the following more general result, which we will prove along the paper:

THEOREM 3. *If $f \in \text{Diff}^1(M)$ is Anosov, if F denotes its unstable sub-bundle, and if μ is an ergodic probability measure for f , then the ϵ -pseudo basin $A_{\epsilon,n}(\mu)$ of μ up to time n satisfies the following equality:*

$$(1.6) \quad \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon,n}(\mu))}{n} = h_\mu(f) - \int \log |\det Df|_F d\mu.$$

1.2. Organization of the paper. In Section 2 we prove Theorem 1, which states the general properties of weak pseudo physical measures for any continuous map $f : M \mapsto M$.

In Section 3, for Anosov diffeomorphisms, we prove part A) of Theorem 2 and also the first part of B). Precisely, we prove that the weak pseudo-physical property of the ergodic components is a sufficient condition to satisfy Pesin's Entropy Formula.

In Section 4, for Anosov diffeomorphisms, we prove the converse statement in part B) of Theorem 2. Namely, the weak pseudo-physical property of the ergodic components is also a necessary condition to satisfy Pesin's Entropy Formula.

Through the proof of Theorem 2, we obtain some stronger intermediate results that hold for any ergodic measure. Finally, at the end of Section 4, we join those intermediate results to prove Theorem 3.

2. Properties of the weak pseudo-physical measures

The purpose of this section is to prove Theorem 1. Along this section, we assume that f is only a continuous map from a compact Riemannian manifold M into itself.

Let us divide the proof of Theorem 1 into its assertions A) to F):

THEOREM 1 A) *Any weak pseudo-physical measure μ is f -invariant.*

PROOF. From Equality (1.4), for any fixed value of $\epsilon > 0$ there exists $n_j \rightarrow +\infty$ such that $\text{Leb}(A_{\epsilon, n_j}(\mu)) > 0$. Thus, there exists $x_j \in M$ such that

$$(2.1) \quad \text{dist}^*(\sigma_{n_j}(x_j), \mu) < \epsilon.$$

Since $\sigma_{n_j}(x_j) \in \mathcal{M}$ and \mathcal{M} is sequentially compact, it is not restrictive to assume that $\{\sigma_{n_j}(x_j)\}$ is weak* convergent. Denote by ν its limit. We assert that ν is f -invariant. In fact, consider the operator $f^* : \mathcal{M} \mapsto \mathcal{M}$ defined by $f^*(\nu)(B) = \nu(f^{-1}(B))$ for any Borel measurable set $B \subset M$. Then $f^*(\delta_y) = \delta_{f(y)}$ for all $y \in M$; hence $f^*(\sigma_{n_j}(x_j)) = \sigma_{n_j}(f(x_j))$ for all $j \in \mathbb{N}$.

It is well known that f^* is continuous. Thus, taking limit in the weak* topology, we obtain:

$$f^*(\nu) = \lim_{j \rightarrow +\infty} f^*(\sigma_{n_j}(x_j)) = \lim_{j \rightarrow +\infty} \sigma_{n_j}(f(x_j)).$$

Since

$$\sigma_{n_j}(x_j) = \frac{1}{n_j} \sum_{i=1}^{n_j-1} \delta_{f^i(x_j)}, \quad \sigma_{n_j}(f(x_j)) = \frac{1}{n_j} \sum_{i=1}^{n_j-1} \delta_{f^{i+1}(x_j)},$$

we deduce that the total variation of the signed measure $\sigma_{n_j}(x_j) - f^*(\sigma_{n_j}(x_j))$ is

$$|\sigma_{n_j}(x_j) - f^*(\sigma_{n_j}(x_j))| \leq \frac{1}{n_j} (\delta_{x_j} + \delta_{f^{n_j}(x_j)}).$$

Thus $\lim_{j \rightarrow +\infty} \sigma_{n_j}(x_j) = \lim_{j \rightarrow +\infty} f^*(\sigma_{n_j}(x_j))$, hence $\nu = f^*(\nu)$, or equivalently ν is f -invariant.

From (2.1) $\text{dist}^*(\nu, \mu) \leq \epsilon$. We have proved that for all $\epsilon > 0$ there exists $\nu \in \mathcal{M}_f$ such that $\text{dist}^*(\mu, \nu) \leq \epsilon$. Since \mathcal{M}_f is sequentially compact, we deduce that $\mu \in \mathcal{M}_f$, as wanted. \square

THEOREM 1 B) *Any physical or pseudo-physical measure is weak pseudo-physical.*

PROOF. Trivially any physical measure is pseudo-physical. So, it is only left to prove that any pseudo-physical measure μ is weak pseudo-physical. Consider $x \in A_\epsilon(\mu)$. From equality (1.2), there exists $n_j \rightarrow +\infty$ such that $\text{dist}^*(\sigma_{n_j}(x), \mu) < \epsilon$.

Therefore, from (1.3) $x \in \bigcap_{N \geq 1} \bigcup_{n \geq N} A_{\epsilon, n}(\mu)$. Since the latter assertions holds for all $x \in A_{\epsilon}(\mu)$, we have proved that

$$A_{\epsilon}(\mu) \subset \bigcap_{N \geq 1} \bigcup_{n \geq N} A_{\epsilon, n}(\mu).$$

As μ is pseudo-physical, we deduce:

$$(2.2) \quad \text{Leb} \left(\bigcap_{N \geq 1} \bigcup_{n \geq N} A_{\epsilon, n}(\mu) \right) \geq \text{Leb}(A_{\epsilon}(\mu)) > 0 \quad \forall \epsilon > 0.$$

Now, assume by contradiction, that μ is not weak pseudo-physical. Taking into account that $\text{Leb}(A_{\epsilon, n}(\mu)) \leq 1$, from the contrary of equality (1.4), we deduce that there exist $\epsilon > 0$ and $a > 0$ such that

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb}(A_{\epsilon, n}(\mu)) = -2a < 0.$$

Therefore, there exists $N \geq 0$ such that $\text{Leb}(A_{\epsilon, n}(\mu)) \leq e^{-an}$ for all $n \geq N$, from

where we deduce that $\sum_{n=1}^{+\infty} \text{Leb}(A_{\epsilon, n}(\mu)) < +\infty$. Finally, applying Borell-Cantelli

Lemma, we conclude that $\text{Leb} \left(\bigcap_{N \geq 1} \bigcup_{n \geq N} A_{\epsilon, n}(\mu) \right) = 0$, contradicting inequality (2.2). \square

THEOREM 1 C) *Weak pseudo-physical measures do exist.*

PROOF. In Theorem 1.3 of [CE], it is proved for any continuous map f on a compact manifold, that the pseudo-physical measures (which in that paper are also called SRB-like or observable) do exist. Since any pseudo-physical measure is weak pseudo-physical, these latter measures always exist. \square

THEOREM 1 D) *The set \mathcal{P}_f of weak pseudo-physical measures does not depend on the choice of the metric in \mathcal{M} that induces the weak* topology.*

PROOF. Take two metrics dist_1^* and dist_2^* , both inducing the weak* topology on \mathcal{M} . We assume that μ is weak pseudo-physical according to dist_1^* , and let us prove that it is also weak pseudo-physical according to dist_2^* .

Since both metric induce the same topology, for any $\epsilon > 0$ there exists $\delta > 0$ such that

$$(2.3) \quad \rho \in \mathcal{M}, \quad \text{dist}_1^*(\rho, \mu) < \delta \quad \Rightarrow \quad \text{dist}_2^*(\rho, \mu) < \epsilon.$$

In the notation of equality (1.3), add a subindex 1 or 2 to denote the sets $A_{\epsilon, n, 1}(\mu)$ and $A_{\epsilon, n, 2}(\mu)$, according to which metric (dist_1^* and dist_2^* , respectively) is used to define them. So, from assertion (2.3) we have: $A_{\delta, n, 1}(\mu) \subset A_{\epsilon, n, 2}(\mu)$, from where

$$\text{Leb}(A_{\delta, n, 1}(\mu)) \leq \text{Leb}(A_{\epsilon, n, 2}(\mu)).$$

Since we are assuming that μ is weak pseudo-physical according to dist_1^* , from equality (1.4) we know that

$$\limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\delta, n, 1}(\mu))}{n} = 0.$$

Then,

$$\limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n, 2}(\mu))}{n} \geq 0.$$

As $\epsilon > 0$ was arbitrarily chosen, the latter inequality holds for all $\epsilon > 0$. But the limit in the latter inequality is non positive because Leb is a probability measure. We conclude that

$$\limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n, 2}(\mu))}{n} = 0 \quad \forall \epsilon > 0,$$

ending the proof that μ is also weak pseudo-physical with respect to the metric dist_2^* . \square

THEOREM 1 E) *The set \mathcal{P}_f of weak pseudo-physical measures is weak*-compact.*

PROOF. Since $\mathcal{P}_f \subset \mathcal{M}$ and \mathcal{M} is weak*-compact, it is enough to prove that \mathcal{P}_f is weak*-closed. Assume $\mu_j \in \mathcal{P}_f$ and $\mu \in \mathcal{M}$ such that

$$\lim_{j \rightarrow +\infty} \text{dist}^*(\mu_j, \mu) = 0$$

We will prove that $\mu \in \mathcal{P}_f$. For any given $\epsilon > 0$, choose and fix j such that $\text{dist}^*(\mu_j, \mu) < \epsilon/2$. Thus, from equality (1.3) and the triangle property, we obtain: $A_{\epsilon/2, n}(\mu_j) \subset A_{\epsilon, n}(\mu)$, from where

$$(2.4) \quad \text{Leb}(A_{\epsilon/2, n}(\mu_j)) \leq \text{Leb}(A_{\epsilon, n}(\mu)).$$

Since $\mu_j \in \mathcal{P}_f$, we can apply equality (1.4) to $\text{Leb}(A_{\epsilon/2, n}(\mu_j))$, which joint with inequality (2.4) implies:

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} \log \text{Leb}(A_{\epsilon, n}(\mu)) \geq 0.$$

Finally, since Leb is a probability measure, we deduce that the above limsup equals 0, concluding that $\mu \in \mathcal{P}_f$ as wanted. \square

THEOREM 1 F) $\lim_{n \rightarrow +\infty} \text{dist}^*(\sigma_n(x), \mathcal{P}_f) = 0$ for Lebesgue almost all $x \in M$.

PROOF. Theorem 1.5 of [CE], states that the distance between $\sigma_n(x)$ and the set of pseudo-physical measures converges to zero with $n \rightarrow +\infty$ for Lebesgue almost all $x \in M$. Since the pseudo-physical measures are contained in \mathcal{P}_f , we trivially deduce the wanted equality. \square

THEOREM 1 G) *If the weak pseudo-physical measure μ is unique, then it is physical and its basin of attraction $B(\mu)$ covers Lebesgue a.e. $x \in M$.*

PROOF. It is an immediate consequence of part F). \square

3. Sufficient condition for Pesin's Entropy Formula

In the sequel we assume that the map $f \in \text{Diff}^1(M)$ is Anosov. The purpose of this section is to deduce, as an immediate consequence from previous known results, part A) of Theorem 2, and the sufficient condition to satisfy Pesin's Entropy Formula in part B) of Theorem 2. Namely, we will deduce that if all the ergodic components of an f -invariant measure μ are weak pseudo-physical, then μ satisfies Pesin's Entropy Formula.

Recall Definition 1.2, which defines the ϵ -pseudo basin $A_{\epsilon,n}(\mu)$ up to time n of a probability measure μ . We will apply the following result:

THEOREM 3.1. [CCE] *Let M be a compact Riemannian manifold of finite dimension. Let $f \in \text{Diff}^1(M)$ be Anosov with hyperbolic splitting $TM = E \oplus F$, where E and F are the stable and unstable sub-bundles respectively. Then, the following inequality holds for any f -invariant $\mu \in \mathcal{M}$:*

$$(3.1) \quad \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon,n}(\mu))}{n} \leq h_\mu(f) - \int \log |\det Df|_F d\mu.$$

PROOF. See Proposition 2.1 in [CCE].

REMARK 3.2. For the non negative Lyapunov exponents, we adopt the notation $\chi_i^+(x)$ as in Definition 1.7. For any $f \in \text{Diff}^1(M)$, Margulis and Ruelle inequality [Mar, R2] states:

$$(3.2) \quad h_\mu(f) \leq \int \sum_{i=1}^{\dim(M)} \chi_i^+(x) d\mu.$$

Thus, Pesin's Entropy Formula holds for an invariant measure μ , if and only if the following inequality holds:

$$h_\mu(f) \geq \int \sum_{i=1}^{\dim(M)} \chi_i^+(x) d\mu.$$

Besides, from Definition 1.5, and from the formula of the integral of the volume form along the unstable sub-bundle F , we obtain the following equality for Anosov diffeomorphisms:

$$\int \sum_{i=1}^{\dim(M)} \chi_i^+(x) d\mu = \int \log |\det Df|_F d\mu.$$

Joining the above assertions, we conclude:

Let $f \in \text{Diff}^1(M)$ be Anosov, and F be its unstable sub-bundle. Then, any f -invariant probability measure μ satisfies Pesin's Entropy Formula if and only if

$$(3.3) \quad h_\mu(f) \geq \int \log |\det Df|_F d\mu.$$

We are ready to deduce part A) of Theorem 2, which is indeed a corollary of Theorem 3.1:

PART A) OF THEOREM 2: *If $f \in \text{Diff}^1(M)$ is Anosov and if μ is a weak pseudo-physical f -invariant measure, then μ satisfies Pesin's Entropy Formula. Therefore, the set of invariant probability measures that satisfy Pesin's Entropy Formula is nonempty.*

PROOF. By contradiction, assume that μ does not satisfy Pesin's Entropy Formula. According to Remark 3.2, inequality (3.3) does not hold:

$$h_\mu(f) - \int \log |\det Df|_F| d\mu < 0.$$

Therefore, applying inequality (3.1) of Theorem 3.1, we conclude that there exists $\epsilon > 0$ such that

$$\limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n}(\mu))}{n} < 0.$$

So, equality (1.4) does not hold; hence μ is not weak pseudo-physical, contradicting the hypothesis. We have proved that all the weak pseudo-physical measures for f satisfy Pesin's Entropy Formula. From part c) of Theorem 1, weak pseudo-physical measures do exist. So, the set of measures that satisfy Pesin's Entropy Formula is nonempty. \square

We now recall the following well known result (see for instance Theorems 4.3.7 and 4.5.6 of [Ke]):

THEOREM 3.3. *Let $f \in \text{Diff}^1(M)$ be Anosov. An f -invariant measure μ satisfies Pesin's Entropy Formula if and only if its ergodic components μ_x satisfy it for μ -a.e. $x \in M$.*

PROOF. On the one hand, we recall that any Anosov C^1 diffeomorphism f is expansive, and for any expansive homeomorphism f on M the metric entropy $h_\mu(f)$ depends upper semi-continuously on the f -invariant measure μ (see for instance Theorem 4.5.6 of [Ke]). So, we can apply the theorem of the Affinity of the Entropy Function (see Theorem 4.3.7 of [Ke]), which states that

$$(3.4) \quad h_\mu(f) = \int h_{\mu_x}(f) d\mu(x),$$

where the measures μ_x for μ -a.e. $x \in M$ are the ergodic components of μ .

On the other hand, the ergodic decomposition theorem states that

$$(3.5) \quad \int \log |\det Df|_F| d\mu = \int d\mu(x) \left(\int \log |\det Df_y|_{F(y)}| d\mu_x(y) \right).$$

Joining Equalities (3.4) and (3.5), and taking into account Margulis and Ruelle inequality (3.2), we deduce that

$$h_\mu(f) - \int \log |\det Df|_F| d\mu = 0$$

if and only if

$$h_{\mu_x}(f) - \int \log |\det Df(y)|_{F(y)}| d\mu_x(y) = 0 \text{ for } \mu\text{-a.e. } x \in M,$$

ending the proof of Theorem 3.3. \square

As a consequence we obtain:

PART B) OF THEOREM 2, SUFFICIENT CONDITION: *If $f \in \text{Diff}^1(M)$ is Anosov and if μ is an invariant measure whose ergodic components μ_x are weak pseudo-physical for μ -a.e. $x \in M$, then μ satisfies Pesin's Entropy Formula.*

PROOF. From the hypothesis, and applying part A) of Theorem 2, we deduce that the ergodic components μ_x of μ satisfy Pesin's Entropy Formula for μ -a.e. $x \in M$. So, from Theorem 3.3, the measure μ also satisfies this formula. \square

4. Necessary condition for Pesin's Entropy Formula

In this section we will prove the necessary condition to satisfy Pesin's Entropy Formula, as stated in part B) of Theorem 2. Precisely, we will prove that if $f \in \text{Diff}^1(M)$ is Anosov, and if the f -invariant measure μ satisfies Pesin's Entropy Formula, then the ergodic components of μ are weak pseudo-physical. We will also prove the equality of Theorem 3 for any ergodic measure μ .

4.1. Previous known properties for Anosov diffeomorphisms.

EXPANSIVITY. Recall that any Anosov diffeomorphism f is expansive (see for instance Lemma 3.4 in [B2]). Namely, there exists a constant $\alpha > 0$, which is called the expansivity constant, such that

$$\text{dist}(f^n(x), f^n(y)) < \alpha \quad \forall n \in \mathbb{Z} \quad \Rightarrow \quad x = y.$$

Given two partitions \mathcal{Q} and \mathcal{R} , the partition $\mathcal{Q} \vee \mathcal{R}$ is defined by

$$\mathcal{Q} \vee \mathcal{R} = \left\{ Q \cap R : Q \in \mathcal{Q}, R \in \mathcal{R} \right\}.$$

METRIC ENTROPY FOR EXPANSIVE SYSTEMS. Recall the following result, which follows from Kolmogorov-Sinai Theorem in the case of expansive homeomorphisms (see for instance, Proposition 2.5 of [B2], or also Theorem 3.2.18 and Lemma 4.5.4 of [Ke]):

If \mathcal{R} is a finite partition whose pieces are Borel measurable sets and have diameter smaller than the expansivity constant α , then $\bigcup_{n=0}^{+\infty} \left\{ \bigvee_{k=-n}^{k=+n} f^{-j} \mathcal{R} \right\}$ generates the Borel σ -algebra, and for any f -invariant measure μ , the metric entropy $h_\mu(f)$ can be computed by:

$$(4.1) \quad h_\mu(f) = \limsup_{n \rightarrow +\infty} \frac{H(\mathcal{R}_n, \mu)}{n}, \quad \text{where } \mathcal{R}_n := \bigvee_{j=0}^{n-1} f^{-j}(\mathcal{R}), \quad \text{and}$$

$$(4.2) \quad H(\mathcal{R}_n, \mu) := - \sum_{Y \in \mathcal{R}_n} \mu(Y) \log(\mu(Y)) \leq \log \# \{Y \in \mathcal{R}_n : \mu(Y) > 0\}.$$

Note: In (4.2) at right, $\#\mathcal{P}$ denotes the number of elements of the finite set \mathcal{P} .

RECTANGLES. Recall the definition of rectangle R in the manifold M for the Anosov diffeomorphism f . (See [B2], page 78.) In particular a rectangle R is proper if $\overline{R} = R = \overline{\text{int}(R)}$. For any proper rectangle R and any $x \in R$ denote

$$W_R^s := \text{connected component}(W^s(x) \cap R) \ni x,$$

$$W_R^u := \text{connected component}(W^u(x) \cap R) \ni x,$$

where $W^s(x)$, $W^u(x)$ are the stable and unstable submanifolds of the point x .

The properties below follow from the definition of rectangle R :

(I) For any pair of points $x, y \in R$ there exists a unique point, which we denote by $[x, y]$, defined by

$$[x, y] \in W_R^s(x) \cap W_R^u(y).$$

(II) There exists a constant $K_R > 0$ such that, if $L \subset R$ is a local embedded C^1 -submanifold, with dimension equal to the unstable dimension, and such that L intersects transversally the local stable manifolds $W_R^s(x)$ for all $x \in R$, then

$$(4.3) \quad \text{Leb}^L(L) \geq K_R^{-1},$$

where Leb^L denotes the Lebesgue measure along L .

We recall the definition of Markov partition $\mathcal{R} = \{R_i\}_{1 \leq i \leq k}$ into rectangles R_i (see [B2], pages 78–79) and the following well known result:

THEOREM 4.1. (EXISTENCE OF MARKOV PARTITIONS).

Let $f \in \text{Diff}^1(M)$ be Anosov. Then, for all $\delta > 0$ there exists a Markov partition whose rectangles have diameter smaller than δ .

PROOF. See Theorem 3.12 of [B2].

DEFINITION 4.2. (DYNAMICAL RECTANGLE)

Let \mathcal{R} be a Markov partition of the manifold M , let $x \in M$ and denote by $R(x)$ the rectangle of \mathcal{R} that contains x . Let n be a positive natural number. The *dynamical rectangle* $R_n(x)$ that contains x is defined by

$$R_n(x) := \bigcap_{j=0}^{n-1} f^{-j}(R(f^j(x))).$$

The following property follows from the definition of Markov Partition (see Condition (b) in [B2], page 79):

$$(4.4) \quad \forall n \geq 0, \quad \text{if } y \in R_n(x) \quad \text{then} \quad W_R^s(y) \subset R_n(x).$$

4.2. Technical Lemmas.

LEMMA 4.3. *Let f be an Anosov diffeomorphism on a compact manifold. Then, for any finite Markov partition \mathcal{R} , there exists a constant $K_0 > 0$ satisfying the following inequality for any f -invariant probability measure μ , for any $0 < \epsilon < 1/4$, for any Borel measurable set $A \subset M$ such that $\mu(A) > 1 - \epsilon$, and for any natural number $n \geq 1$:*

$$(4.5) \quad \log \#\{Y \in \mathcal{R}_n : Y \cap A \neq \emptyset\} \geq H(\mathcal{R}_n, \mu) - n \cdot K_0 \cdot \epsilon + \epsilon \log \epsilon + (1 - \epsilon) \log(1 - \epsilon).$$

PROOF. Denote $A_n := \bigcup \{Y \in \mathcal{R}_n : A \cap Y \neq \emptyset\}$. Since $A \subset A_n$ we have $\mu(A_n) > 1 - \epsilon$. If $\mu(A_n) = 1$ then inequality (4.5) holds trivially as a consequence of (4.2). So, let us prove Lemma 4.3 in the case

$$1 - \epsilon < \mu(A_n) < 1; \quad \text{hence } 0 < \mu(M \setminus A_n) < \epsilon.$$

By definition:

$$\begin{aligned} H(\mathcal{R}_n, \mu) &:= - \sum_{Y \in \mathcal{R}_n} \mu(Y) \log \mu(Y) \\ &= - \sum_{Y \subset A_n} \mu(Y) \log \mu(Y) - \sum_{Y \subset M \setminus A_n} \mu(Y) \log \mu(Y) \\ &= -\mu(A_n) \sum_{Y \subset A_n} \frac{\mu(Y)}{\mu(A_n)} \log \left(\frac{\mu(Y)}{\mu(A_n)} \right) - \sum_{Y \subset A_n} \mu(Y) \log \mu(A_n) \\ &\quad - \mu(M \setminus A_n) \sum_{Y \subset M \setminus A_n} \frac{\mu(Y)}{\mu(M \setminus A_n)} \log \left(\frac{\mu(Y)}{\mu(M \setminus A_n)} \right) - \sum_{Y \subset M \setminus A_n} \mu(Y) \log \mu(M \setminus A_n). \end{aligned}$$

Construct the probability measures μ_1 and μ_2 defined by the following equalities for all Borelian set $B \subset M$:

$$\mu_1(B) := \mu(B \cap A_n)/\mu(A_n), \quad \mu_2(B) := \mu(B \cap (M \setminus A_n))/\mu(M \setminus A_n).$$

We obtain

$$\begin{aligned} H(\mathcal{R}_n, \mu) &= -\mu(A_n) \sum_{Y \subset A_n} \mu_1(Y) \log(\mu_1(Y)) - \mu(A_n) \log \mu(A_n) \\ &\quad - \mu(M \setminus A_n) \sum_{Y \subset M \setminus A_n} \mu_2(Y) \log(\mu_2(Y)) - \mu(M \setminus A_n) \log \mu(M \setminus A_n). \end{aligned}$$

Applying inequality (4.2):

$$\begin{aligned} H(\mathcal{R}_n, \mu) &\leq \log \#\{Y \in \mathcal{R}_n : Y \subset A_n\} - \mu(A_n) \log \mu(A_n) \\ &\quad + \mu(M \setminus A_n) \cdot \log \#\mathcal{R}_n - \mu(M \setminus A_n) \log \mu(M \setminus A_n). \end{aligned}$$

Taking into account that $0 < \mu(M \setminus A_n) < \epsilon < 1/4$ and that $-u \log u$ is strictly increasing for $0 < u < 1/4$ and strictly decreasing for $u > 3/4$, we obtain:

$$\begin{aligned} H(\mathcal{R}_n, \mu) &\leq \\ \log \#\{Y \in \mathcal{R}_n : Y \subset A_n\} &+ \mu(M \setminus A_n) \cdot \log \#\mathcal{R}_n - \epsilon \log \epsilon - (1 - \epsilon) \log(1 - \epsilon) \leq \\ \log \#\{Y \in \mathcal{R}_n : Y \subset A_n\} &+ \epsilon \cdot n \cdot K_0 - \epsilon \log \epsilon - (1 - \epsilon) \log(1 - \epsilon), \end{aligned}$$

where

$$K_0 := \sup_{n \geq 1} \frac{\log \#\mathcal{R}_n}{n} > 0$$

Therefore, to end the proof of Lemma 4.3 it is enough to show that $K_0 < +\infty$. In fact, for a Markov partition \mathcal{R} , any rectangle $Y \in \mathcal{R}_n$ is obtained as a connected component of the intersection $f^{-n}(R_i) \cap R_j$ for some pair of rectangles $R_i, R_j \in \mathcal{R}$. Fixing $R_i \in \mathcal{R}$, the maximum number of connected components of the intersections of $f^{-n}(R_i)$ with the rectangle R_j of the partition, is upper bounded the following quotient

$$\begin{aligned} &\frac{\max \left\{ \text{Leb}(W_{f^{-n}(R_i)}^s(y)) : y \in f^{-n}(R_i) \right\}}{\min \left\{ \text{Leb}(W_{R_j}^s(x)) : x \in R_j \right\}} \leq \\ &d^n \frac{\max \left\{ \text{Leb}(W_{R_i}^s(x)) : x \in R_i \right\}}{\min \left\{ \text{Leb}(W_{R_j}^s(x)) : x \in R_j \right\}} =: d^n \cdot q_{i,j}, \end{aligned}$$

where

$$d := \max \left\{ |\det Df_x^{-1}|_{E(x)}| : x \in M \right\}.$$

Thus, denoting $k := \#\mathcal{R}$, we have

$$\#\mathcal{R}_n \leq k^2 \cdot d^n \cdot \max \left\{ q_{i,j} : 1 \leq i, j \leq k \right\}.$$

We conclude that

$$\limsup_{n \rightarrow +\infty} \frac{\log \#\mathcal{R}_n}{n} \leq d < +\infty,$$

which implies $K_0 := \sup_{n \geq 1} \frac{\log \#\mathcal{R}_n}{n} < +\infty$, ending the proof of Lemma 4.3. \square

In the following Lemma we will construct a local C^1 -foliation, whose leaves are *pseudo-unstable manifolds* ϵ -approaching (in the C^1 -topology) the true local unstable manifolds of any rectangle of a given Markov partition.

LEMMA 4.4. *Let $f \in \text{Diff}^1(M)$ be Anosov. Denote the stable and unstable subbundles by E and F , respectively. Denote the expansivity constant by $\alpha > 0$. Then, for all $\epsilon > 0$ there exist $0 < \delta_0 < \alpha$ and $K > 0$ such that, for any finite Markov partition $\mathcal{R} = \{R_i\}_{1 \leq i \leq k}$ into rectangles with diameter smaller than δ_0 , there exists a finite family $\{\mathcal{L}_i\}_{1 \leq i \leq k}$ of local foliations \mathcal{L}_i , each one defined in an open neighborhood of each rectangle R_i , satisfying the following properties for all $1 \leq i \leq k$, for all $x \in R_i$ and for all $n \geq 0$:*

A) \mathcal{L}_i is C^1 -trivializable and its leaves are $\dim(F)$ -dimensional.
 B) $\text{dist}(F_{f^n(x)}, T_{f^n(x)} f^n(\mathcal{L}_i(x))) < \epsilon$, where $\mathcal{L}_i(x)$ denotes the leaf of the foliation \mathcal{L}_i that contains x .

C) $K^{-1} e^{-n\epsilon} \leq \frac{|\det dDf_x^n|_{T_x(\mathcal{L}_i(x))}|}{|\det Df_x^n|_{F(x)}} \leq K e^{n\epsilon}$.

D) There exist a point $x_i \in R_i$ and an open subset $A_i^s \subset W_{R_i}^s(x_i)$, in the topology of the stable submanifold $W^s(x_i)$, such that

$$\text{Leb}^{W^s(x_i)}(A_i^s) \geq K^{-1},$$

where $\text{Leb}^{W^s(x_i)}$ denotes the Lebesgue measure along the submanifold $W^s(x_i)$; and besides, if $y \in A_i^s$, then

$$\text{Leb}^{f^n(\mathcal{L}_i(y))}(f^n(\mathcal{L}_i(y) \cap R_n(x))) \geq K^{-1},$$

where $\text{Leb}^{f^n(\mathcal{L}_i(y))}$ denotes the Lebesgue measure along the submanifold $\mathcal{L}_i(y)$.

PROOF. Proposition 3.6 of [CCE] states the existence of $\delta_0 > 0$ and the local C^1 -foliation \mathcal{L}_i satisfying (A), (B) and (C). So, it is enough to prove that if $\epsilon > 0$ is small enough, then any local C^1 -foliation \mathcal{L}_i defined in a neighborhood of the rectangle R_i and satisfying (A) and (B), also satisfies (D) for some constant $K > 0$.

In fact, choose and fix any point x_i in the interior of the rectangle R_i . From the definition of rectangle, for each $z \in R_i$ there exists a unique point in the transversal intersection

$$W_{R_i}^s(z) \cap W_{R_i}^u(x_i) \neq \emptyset.$$

By continuity of the transversal intersection between C^1 -manifolds, there exists $\epsilon' > 0$ such that the following assertion holds:

If $\text{dist}(x_i, y) < \epsilon'$ and if \mathcal{L}_i is any local foliation whose leaves have dimension $\dim(F)$, are C^1 , and are ϵ' -near the unstable local leaves of R_i in the C^1 -topology, then for each $z \in R_i$ the intersection $W_{R_i}^s(z) \cap \mathcal{L}_i(y)$ is transversal and contains a single point.

In particular, we obtain:

$$(4.6) \quad W_{R_i}^s(z) \cap \mathcal{L}_i(y) \neq \emptyset$$

$$\forall y \in W_{R_i}^s(x_i) \text{ such that } \text{dist}(x_i, y) < \epsilon', \quad \forall z \in R_n(x) \subset R_i, \quad \forall x \in R_i.$$

Define

$$A_i^s := \left\{ y \in W_{R_i}^s(x_i) : \text{dist}(x_i, y) < \epsilon' \right\}.$$

By construction A_i^s is an open subset of the local stable submanifold $W_{R_i}^s(x_i)$, in the topology of this submanifold. Construct a real number $K_i > 0$ large enough so

$$(4.7) \quad \text{Leb}^{W^s(x_i)}(A_i^s) \geq K_i^{-1}.$$

From assertion (4.6) we deduce

$$f^n(\mathcal{L}_i(y) \cap R_n(x)) \cap W_{f^n(R_n(x))}^s(w) \neq \emptyset$$

for all $y \in A_i^s$ and for all $w \in f^n(R_n(x))$.

From the definition of the dynamical rectangle $R_n(x)$ and from the properties of the Markov partition, there exists a rectangle $R_j \ni f^n(x)$ of the partition such that local stable manifold $W_{R_j}^s(w) \supset W_{f^n(R_n(x))}^s(w)$ for all $w \in R_j$.

So, we deduce

$$f^n(\mathcal{L}_i(y) \cap R_n(x)) \cap W_{R_j}^s(w) \neq \emptyset$$

for all $y \in A_i^s$ and for all $w \in R_j$. In other words, the pseudo-unstable $\dim(F)$ -submanifold $f^n(\mathcal{L}_i(y) \cap R_n(x))$ intersects transversally all the local stable submanifolds of the rectangle R_j where it is contained. Thus, applying inequality (4.3) we have

$$(4.8) \quad \text{Leb}^{f^n(\mathcal{L}_i)} f^n(\mathcal{L}_i(y) \cap R_n(x)) \geq \frac{1}{K_{R_j}}.$$

Finally, define $K = \max_{1 \leq i \leq k} \{K_{R_i}, K_i\}$. From inequalities (4.7) and (4.8), we conclude assertion D), as wanted. \square

4.3. End of the proofs of Theorems 2 and 3. For any probability measure μ recall equality (1.3), defining the measurable set $A_{\epsilon, n}(\mu)$ which we called the ϵ -pseudo basin of μ up to time n . We will end the proof of Theorem 2, by applying the following key result which bounds from below the Lebesgue measure of the set $A_{\epsilon, n}(\mu)$ for any ergodic measure μ :

THEOREM 4.5. *Let M be a compact Riemannian manifold of finite dimension. Let $f \in \text{Diff}^1(M)$ be Anosov with hyperbolic splitting $TM = E \oplus F$, where E and F are the stable and unstable sub-bundles respectively. Let μ be an ergodic measure. Then:*

$$(4.9) \quad \lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n}(\mu))}{n} \geq h_\mu(f) - \int \log |\det Df|_F d\mu.$$

PROOF. We notice that the limit at left in equality (4.9) does not depend on the choice of metric dist^* that induces the weak* topology in the space \mathcal{M} of Borel probability measures. In fact, to prove the latter assertion it is enough to argue as in the proof of part (D) of Theorem 1 in Section 2. So, to prove Theorem 4.5 we choose and fix the following metric in \mathcal{M} :

$$\text{dist}^*(\mu, \nu) := \sum_{i=0}^{+\infty} \frac{|\int \varphi_i d\mu - \int \varphi_i d\nu|}{2^i} \quad \forall \mu, \nu \in \mathcal{M},$$

where $\{\varphi_i\}_{i \in \mathbb{N}}$ is any fixed countable family of real continuous functions $\varphi_i \in C^0(M, [0, 1])$ that is dense in $C^0(M, [0, 1])$. Note that, according to the metric dist^* , the balls are convex. In other words, if a finite number of probability measures belong to the ball with centre μ and radius $\epsilon > 0$, then any convex combination of those measures also belongs to it.

For any $x \in M$ and for any natural number $n \geq 1$ denote:

$$\begin{aligned}\psi(x) &:= \log |\det(Df_x|_{F(x)})|, \\ \psi_n(x) &:= \log |\det(Df_x^n|_{F(x)})| = \sum_{j=0}^{n-1} \psi(f^j(x)) = n \cdot \int \psi d\sigma_n(x),\end{aligned}$$

where $\sigma_n(x)$ is the empiric probability constructed in Definition 1.1.

Fix any real value $\epsilon > 0$. The real function $\psi : M \mapsto \mathbb{R}$ is continuous because f is of class C^1 and the sub-bundle F is continuous. Thus, from the definition of the weak* topology in the space \mathcal{M} of probability measures, we deduce that there exists $0 < \epsilon' < \epsilon$ such that

$$\nu, \mu \in \mathcal{M}, \quad \text{dist}^*(\nu, \mu) < \epsilon' \quad \Rightarrow \quad \left| \int \psi d\nu - \int \psi d\mu \right| < \epsilon.$$

In particular, for $\nu = \sigma_n(x)$ we deduce:

$$(4.10) \quad \text{If } \text{dist}^*(\sigma_n(x), \mu) < \epsilon', \text{ then } \left| \log |\det(Df_x^n|_{F(x)})| - n \cdot \int \psi d\mu \right| < n \cdot \epsilon.$$

Since μ is an ergodic probability measure, we have $\lim_{n \rightarrow +\infty} \sigma_n(x) = \mu$ for μ -a.e. $x \in M$. So, for the fixed value of $\epsilon' > 0$ as above, and for μ -a.e. $x \in M$, there exists $N(x) \geq 1$ such that

$$\text{dist}^*(\sigma_n(x), \mu) < \epsilon'/2 \quad \forall n \geq N(x).$$

For any natural value of $N \geq 1$, define the set

$$(4.11) \quad A_N := \left\{ x \in M : \text{dist}^*(\sigma_n(x), \mu) < \epsilon'/2 \quad \forall n \geq N \right\}.$$

Since $A_N \subset A_{N+1}$ and $\mu\left(\bigcup A_N\right) = 1$, there exists $N \geq 1$ such that

$$(4.12) \quad \mu(A_N) \geq 1 - \epsilon.$$

In the sequel, we fix such a value of $N \geq 1$.

From the definition of the metrizable weak*-topology in the space \mathcal{M} of Borel-probability measures, it is standard to check that the Dirac delta probability δ_x depends uniformly continuously on the point $x \in M$. Since the empiric probability $\sigma_n(x)$ is a convex combination of Dirac delta measures, and the balls in \mathcal{M} are convex, we deduce that there exists $\delta_2 > 0$ such that, for any pair of points $x, y \in M$ and for any natural value of $n \geq 1$, the following assertion holds:

$$(4.13) \quad \text{If } \text{dist}(f^j(x), f^j(y)) < \delta_2 \text{ for all } 0 \leq j \leq n-1, \text{ then } \text{dist}^*(\sigma_n(x), \sigma_n(y)) < \epsilon'/2.$$

For the fixed value of $\epsilon > 0$ at the beginning, we construct the real numbers $0 < \delta_0 < \alpha$ (where α is expansivity constant), and $K > 0$, as in Lemma 4.4. We consider any Markov partition $\mathcal{R} = \{R_i\}_{1 \leq i \leq k}$ with diameter smaller than $\min\{\delta_0, \delta_1, \delta_2\}$ and, for each rectangle R_i , we construct the C^1 -foliation \mathcal{L}_i that satisfies the properties (A) to (D) of Lemma 4.4.

From equality (4.11), assertion (4.13), and the triangle property of the metric, we deduce the following assertion for all $n \geq N$:

$$\text{If } x \in A_N \text{ and } y \in R_n(x), \text{ then}$$

$$(4.14) \quad \text{dist}^*(\sigma_n(y), \mu) \leq \text{dist}^*(\sigma_n(y), \sigma_n(x)) + \text{dist}^*(\sigma_n(x), \mu) < \frac{\epsilon'}{2} + \frac{\epsilon'}{2} = \epsilon' < \epsilon.$$

Recalling equality (1.3), from the above assertion we deduce that $y \in A_{\epsilon, n}(\mu)$ for all $y \in R_n(x)$. Since the rectangle $R_n(x)$ is any piece of the partition $\mathcal{R}_n = \bigvee_{j=0}^{n-1} \mathcal{R}$ that intersects A_N , we deduce the following statement for all $n \geq N$:

If $Y \in \mathcal{R}_n$ and $Y \cap A_N \neq \emptyset$, then $Y \subset A_{\epsilon, n}(\mu)$.

Therefore

$$(4.15) \quad \text{Leb}(A_{\epsilon, n}(\mu)) \geq \sum_{Y \in \mathcal{R}_n, Y \cap A_N \neq \emptyset} \text{Leb}(Y).$$

Besides, joining assertion (4.10) and inequality (4.14), we deduce the following property for all $n \geq N$:

If $Y \in \mathcal{R}_n$ and $Y \cap A_N \neq \emptyset$, then

$$(4.16) \quad \left| \log |\det(Df_y^n|_{F(y)})| - n \cdot \int \psi d\mu \right| < n \cdot \epsilon \quad \forall y \in Y.$$

Now, for any $n \geq N$, let us compute $\text{Leb}(Y)$ for any rectangle $Y \in \mathcal{R}_n$ such that $Y \cap A_N \neq \emptyset$. Since $Y \subset R_i \in \mathcal{R}$, to compute $\text{Leb}(Y)$ we will use the Fubini decomposition of the Lebesgue measure along the local pseudo-unstable C^1 -foliation \mathcal{L}_i . Applying part (D) of Lemma 4.4 consider the point $x_i \in R_i$ and the submanifold $A_i^s \subset W_{R_i}^s(x_i)$. Taking the Fubini decomposition of Leb we obtain:

$$\text{Leb}(Y) = \int_{z \in W_{R_i}^s(x_i)} d\text{Leb}^{W^s(x_i)}(z) \int_{y \in \mathcal{L}_i(z) \cap Y} |\det D\phi_i(y)| d\text{Leb}^{\mathcal{L}_i(z)}(y),$$

where ϕ_i^{-1} is a local C^1 -diffeomorphism that parameterizes the neighborhood of R_i and trivializes the C^1 -foliation \mathcal{L}_i . Therefore, $|\det D\phi_i|$ is continuous and bounded away from zero by a constant, say $k_i > 0$. Since A_i^s is an open subset of $W_{R_i}^s(x_i)$ in the topology of this local stable manifold, we obtain:

$$\text{Leb}(Y) \geq k_i \cdot \int_{z \in A_i^s} d\text{Leb}^{W^s(x_i)}(z) \int_{y \in \mathcal{L}_i(z) \cap Y} d\text{Leb}^{\mathcal{L}_i(z)}(y).$$

Changing variables $y' = f^n(y)$ in the integral at right, we obtain:

$$\text{Leb}(Y) \geq k_i \cdot \int_{z \in A_i^s} I(z) d\text{Leb}^{W^s(x_i)}(z), \text{ where}$$

$$I(z) = \int_{y' \in f^n(\mathcal{L}_i(z) \cap Y)} \left| \det Df_{f^{-n}(y')}^n|_{T_{f^{-n}(y')} \mathcal{L}_i(z)} \right|^{-1} d\text{Leb}^{f^n(\mathcal{L}_i(z))}(y').$$

Since $y \in Y \subset R_i$, we can apply inequality at left of part c) of Lemma 4.4:

$$\text{Leb}(Y) \geq k_i \cdot K^{-1} \cdot e^{-n\epsilon} \cdot \int_{z \in A_i^s} J(z) d\text{Leb}^{W^s(x_i)}(z), \text{ where}$$

$$J(z) = \int_{y' \in f^n(\mathcal{L}_i(z) \cap Y)} \left| \det Df_{f^{-n}(y')}^n|_{F(f^{-n}(y'))} \right|^{-1} d\text{Leb}^{f^n(\mathcal{L}_i(z))}(y').$$

Since $y = f^{-n}(y') \in Y$ and $Y \cap A_N \neq \emptyset$, we can apply inequality (4.16):

$$\text{Leb}(Y) \geq k_i \cdot K^{-1} \cdot e^{-2n\epsilon - n \int \psi d\mu} \cdot J, \text{ where}$$

$$J = \int_{z \in A_i^s} d\text{Leb}^{W^s(x_i)}(z) \int_{y' \in f^n(\mathcal{L}_i(z) \cap Y)} d\text{Leb}^{f^n(\mathcal{L}_i(z))}(y') =$$

$$\int_{z \in A_i^s} \text{Leb}^{f^n(\mathcal{L}_i(z))}(f^n(\mathcal{L}_i(z))) d\text{Leb}^{W^s(x_i)}(z).$$

From part (D) of Lemma 4.4 we know that $\text{Leb}^{f^n(\mathcal{L}_i(z))}(f^n(\mathcal{L}_i(z))) \geq K^{-1}$ for all $z \in A_i^s$, and besides $\text{Leb}^{W^s(x_i)}(A_i^s) \geq K^{-1}$. Thus, we have proved the following inequality for all $n \geq N$, and for all $Y \in \mathcal{R}_n$ such that $Y \cap A_N \neq \emptyset$:

$$\text{Leb}(Y) \geq k_i \cdot K^{-3} \cdot e^{-2n\epsilon - n \int \psi d\mu}.$$

Joining the above inequality with inequality (4.15), we deduce, for all $n \geq N$:

$$\text{Leb}(A_{\epsilon, n}(\mu)) \geq k_i \cdot K^{-3} \cdot e^{-2n\epsilon - n \int \psi d\mu + \log \#\{Y \in \mathcal{R}_n : Y \cap A_N \neq \emptyset\}}.$$

Therefore,

$$\limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n}(\mu))}{n} \geq$$

$$-2\epsilon - \int |\det Df|_F d\mu + \limsup_{n \rightarrow +\infty} \frac{\log \#\{Y \in \mathcal{R}_n : Y \cap A_N \neq \emptyset\}}{n}.$$

Finally, applying Lemma 4.3, we deduce that

$$\limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n}(\mu))}{n} \geq$$

$$-2\epsilon - \int |\det Df|_F d\mu + \limsup_{n \rightarrow +\infty} \frac{H(\mathcal{R}_n, \mu)}{n} - K_0 \cdot \epsilon \quad \forall 0 < \epsilon < 1.$$

So, from equality (4.1), we conclude

$$\lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n}(\mu))}{n} \geq h_\mu(f) - \int |\det Df|_F d\mu,$$

ending the proof of Theorem 4.5. \square

Now, we are ready to end the proofs of Theorems 2 and 3, as consequences of Theorems 4.5, 3.3 and 3.1:

PART B) OF THEOREM 2, NECESSARY CONDITION: *If $f \in \text{Diff}^1(M)$ is Anosov and if μ is an invariant measure satisfying Pesin's Entropy Formula, then its ergodic components μ_x are weak pseudo-physical for μ -a.e. $x \in M$.*

PROOF. First, let us assume that μ is ergodic satisfying Pesin's Entropy Formula. From Theorem 4.5 we obtain

$$\lim_{\epsilon \rightarrow 0^+} \limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n}(\mu))}{n} \geq 0.$$

From equality (1.3), if $\epsilon_1 < \epsilon_2$ then $A_{\epsilon_1, n}(\mu) \subset A_{\epsilon_2, n}(\mu)$.

So $\limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n}(\mu))}{n}$ is increasing with $\epsilon > 0$. Thus

$$\limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n}(\mu))}{n} \geq 0 \quad \forall \epsilon > 0.$$

But since Leb is a probability measure, we conclude that

$$\limsup_{n \rightarrow +\infty} \frac{\log \text{Leb}(A_{\epsilon, n}(\mu))}{n} = 0 \quad \forall \epsilon > 0.$$

Applying Definition 1.3, we deduce that μ is weak pseudo-physical.

We have proved that any ergodic measure that satisfies Pesin's Entropy Formula is weak pseudo-physical. Now let us consider a non ergodic measure μ that satisfies Pesin's Entropy Formula. From Theorem 3.3 we know that its ergodic components μ_x also satisfy that formula for μ -a.e. $x \in M$. We conclude that the ergodic components μ_x of μ are weak pseudo-physical for μ -a.e. $x \in M$, as wanted. \square

Finally to complete all the proofs, we add the following immediate end:

END OF THE PROOF OF THEOREM 3.

PROOF. The equality of Theorem 3 is immediately obtained by joining the inequalities of Theorems 3.1 and 4.5. \square

References

- [BDV] C. Bonatti, L. Diaz, M. Viana: *Dynamics Beyond Uniform Hyperbolicity: A Global Geometric and Probabilistic Perspective*, Springer-Verlag, Berlin Heidelberg, 2005.
- [B1] R. Bowen: *Periodic points and measures for Axiom A diffeomorphisms*. Trans. Amer. Math. Soc. 154 (1971), pp. 377–397.
- [B2] R. Bowen: *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms* Lect. Notes in Math. 470 Springer-Verlag, New York, 1975.
- [CCE] E. Catsigeras, M. Cerminara, H. Enrich: *The Pesin Entropy Formula for C^1 Diffeomorphisms with Dominated Splitting*. Ergod.Th.& Dynam.Sys. 35 (2015), pp. 737–761.
- [CE] E. Catsigeras, H. Enrich: *SRB-like measures for C^0 dynamics*, Bull. Pol. Acad. Sci. Math. 59 (2011), pp. 151–164.
- [GI] A. Gorodetski, Yu. S. Ilyashenko: *Minimal and strange attractors*. Journ. Bif. & Chaos 6 (1996), pp. 1177–1183.
- [HPS] M. Hirsch, C. Pugh, M. Shub: *Invariant manifolds* Lect. Notes in Math. 587, Springer-Verlag, Berlin, 1977.
- [II] Yu. S. Ilyashenko: *Minimal attractors*. In *Proceedings of Equadiff 2003*. World Sci. Pub., Singapore, 2005, pp. 421–428.
- [Ke] G. Keller: *Equilibrium States in Ergodic Theory*. London Math. Soc. Student Texts Vol. 42, Cambridge Univ. Press, Cambridge, 1998.
- [LY] F. Ledrappier, L-S. Young: *The metric entropy of diffeomorphisms. Part I: Characterization of measures satisfying Pesin's formula*. Ann. Math. 122 (1985), pp. 509–539.
- [Mañ] R. Mañé: *A proof of Pesin's Formula*. Ergod. Th. & Dynam. Sys. 1, pp. 95–102, 1981. *Errata*. Ergod.Th.& Dynam.Sys. 3 (1983), pp. 159–160.
- [Mar] G.A. Margulis: *Proof of an Inequality for the Metric Entropy* Unpublished, 1966. Cited in Hasselblatt and Pesin: *Pesin's Entropy Formula* Scholarpedia 3 (2008), pp. 3733–3733.
- [P1] Ya. B. Pesin: *Families of invariant manifolds corresponding to nonzero characteristic exponents*. Math USSR-Izv. 40 (1976), pp. 1261–1307.
- [P2] Ya. B. Pesin: *Characteristic Lyapunov Exponents and Smooth Ergodic Theory*. Russian Math. Surveys 32 (1977), pp. 55–114.

- [PS] Ya. B. Pesin, Ya. G. Sinai: *Gibbs measures for partially hyperbolic attractors*. Ergod.Th.& Dynam.Sys. 2 (1982), pp 417–438.
- [Pu] C. Pugh: *The $C^{1+\alpha}$ hypothesis in Pesin theory*. Inst. Hautes Études Sci. Pub. Math. 59 (1984), pp. 143–161.
- [Qiu] H. Qiu: *Existence and uniqueness of SRB measure on C^1 generic hyperbolic attractors*, Commun. Math. Phys. 302 (2011), pp. 345–357.
- [RoY] C. Robinson, L-S. Young: *Nonabsolutely Continuous Foliations for an Anosov Diffeomorphism*. Invent. Math. 61 (1980), pp. 159–176.
- [R1] D. Ruelle: *A measure associated with axiom A attractors*. Amer. Journ. of Math. 98 (1976), pp. 619–654.
- [R2] D. Ruelle: *An inequality for the entropy of differentiable maps*. Bol. Soc. Bras. Mat. 9 (1978), pp. 83–87.
- [Sh] M. Shub: *Dynamical systems, filtrations and entropy*, Bull. Amer. Math. Soc. 80 (1974), pp. 27–41.
- [Si] Ya. G. Sinai: *Gibbs measure in ergodic theory*. Russ. Math. Surv. 27 (1972), pp. 21–69.
- [ST] W. Sun and X. Tian: *Dominated Splitting and Pesin’s Entropy Formula*, Disc. & Cont. Dyn. Sys. 32 (2012), pp. 1421–1434.
- [Ta] A. Tahzibi: *C^1 -generic Pesin’s entropy formula*, C. R. Acad. Sci. Paris, Ser I, 335 (2002), pp. 1057–1062.
- [YC] D. Yang, Y. Cao: *On Pesin’s entropy formula for dominated splittings without mixed behavior*, ArXiv: 1512.08541v1 [math.DS], 2015

INSTITUTO DE MATEMÁTICA Y ESTADÍSTICA “RAFAEL LAGUARDIA”(IMERL), FACULTAD DE INGENIERÍA, UNIVERSIDAD DE LA REPÚBLICA. URUGUAY.

E-mail address: `eleonora@fing.edu.uy`

E-mail address: `cerminar@fing.edu.uy`

E-mail address: `enrich@fing.edu.uy`